

REFERENCE NO.

35

In re application of: Hong Liu et al.
Application No.: not yet assigned
Filing Date: March 23, 2004
Attorney Docket No.: 021288-002610US

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/09110 A1

(51) International Patent Classification⁷: **C07D 295/18**,
235/18, 211/66, 207/16, 403/06, A61K 31/5375, A61P
37/00, C07C 255/31

(21) International Application Number: **PCT/US00/20453**

(22) International Filing Date: **28 July 2000 (28.07.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/146,647 **30 July 1999 (30.07.1999)** **US**

(71) Applicant: **BOEHRINGER INGELHEIM PHARMA-
CEUTICALS, INC.** [US/US]; 900 Ridgebury Road, P.O.
Box 368, Ridgefield, CT 06877 (US).

(72) Inventors: **BEKKALI, Younes**; Unit 34, 60 Padanaram
Road, Danbury, CT 06811 (US). **BETAGERI, Raj**; 62
Redwood Drive, Bethel, CT 06801 (US). **EMMANUEL,
Michel**; 22 B Myrtle Avenue, Danbury, CT 06810 (US).
HICKEY, Eugene; 5 Woodbury Drive, Danbury, CT
06811 (US). **LIU, Weimin**; 56 Wakelee Avenue Ext.-#12,
Shelton, CT 06484 (US). **SPERO, Denice, M.**; 18

Limekiln Road, West Redding, CT 06896 (US). **THOM-
SON, David, S.**; 166 Minuteman Road, Ridgefield, CT
06877 (US). **WARD, Yancey**; 11 Narragansett Trail,
Sandy Hook, CT 06482 (US). **YOUNG, Erick, R., R.**;
10232 Avalon Valley Drive, Danbury, CT 06810 (US).
PATEL, Usha; 20 Dairy Farm Drive, Brookfield, CT
06804 (US).

(74) Agents: **RAYMOND, Robert, P. et al.**; Boehringer In-
gelheim Corporation, 900 Ridgebury Road, P.O. Box 368,
Ridgefield, CT 06877 (US).

(81) Designated States (*national*): **CA, JP, MX.**

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).


Published:

- *With international search report.*
- *Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/09110 A1

(54) Title: **NOVEL SUCCINATE DERIVATIVE COMPOUNDS USEFUL AS CYSTEINE PROTEASE INHIBITORS**



(57) Abstract: Disclosed are novel succinate
derivative compounds of formula (I)/(Ia); wherein
R1, R2, R3, R4, R5, R6, R7, X and A are defined
herein. The compounds are useful as inhibitors of
cysteine proteases. Also disclosed are methods of
using and methods of making such compounds.

Novel Succinate Derivative Compounds Useful as Cysteine Protease InhibitorsRELATED APPLICATION DATA

This application claims benefit to US provisional application no. 60/146,647 filed

5 7/30/99.

TECHNICAL FIELD OF THE INVENTION

This invention relates to peptidyl cysteine protease inhibitors. The compounds are reversible inhibitors of the cysteine proteases S, K, F, L and B and are therefore useful in
10 the treatment of autoimmune and other cathepsin related diseases. The invention also discloses processes for preparing such compounds and pharmaceutical compositions comprising them.

BACKGROUND OF THE INVENTION

15 Cathepsin S and cathepsin K are members of the papain family, within the papain superfamily of cysteine proteases. The papain family is the largest group of cysteine proteases and includes proteases such as cathepsins B, H, K, L, O and S. (A.J. Barrett et al., 1996, Perspectives in Drug Discovery and Design, 6, 1). The cysteine proteases have important roles in human biology and diseases including atherosclerosis, emphysema,
20 osteoporosis, chronic inflammation and immune disorders (H.A. Chapman et al., 1997, Ann. Rev. Physiol., 59, 63). Cathepsin S plays a key role in regulating antigen presentation and immunity (H.A. Chapman, 1998, Current Opinion in Immunology, 10, 93; R. J. Riese et al., 1998, J. Clin. Invest., 101, 2351; R.J. Riese et al., 1996, Immunity, 4, 357). Cathepsin S deficient mice have impaired invariant chain degradation resulting in
25 decreased antigen presentation and germinal center formation, and diminished susceptibility to collagen-induced arthritis indicating the therapeutic potential for a cathepsin S inhibitor (G. Shi et al., 1999, Immunity, 10, 197; T.Y. Nakagawa et al, 1999, Immunity, 10, 207) .

30 The specificity of the immune response relies on processing of foreign protein and presentation of antigenic peptide at the cell surface. Antigenic peptide is presented bound

to MHC Class II, a heterodimeric glycoprotein expressed in certain antigen presenting cells of hematopoietic lineage, such as B cells, macrophages and dendritic cells.

Presentation of antigen to effector cells, such as T-cells, is a fundamental step in recognition of non-self and thus initiation of the immune response.

5

Recently MHC Class II heterodimers were shown to associate intracellularly with a third molecule designated invariant chain. Invariant chain facilitates Class II transport to the endosomal compartment and stabilizes the Class II protein prior to loading with antigen. Invariant chain interacts directly with Class II dimers in the antigen-binding groove and
10 therefore must be proteolyzed and removed or antigen cannot be loaded or presented. Current research suggests that invariant chain is selectively proteolyzed by cathepsin S, which is compartmentalized with MHC Class II complexes within the cell. Cathepsin S degrades invariant chain to a small peptide, termed CLIP, which occupies the antigen – binding groove. CLIP is released from MHC Class II by the interaction of MHC Class II
15 with HLA-DM, a MHC-like molecule thus freeing MHC Class II to associate with antigenic peptides. MHC Class II-antigen complexes are then transported to the cell surface for presentation to T-cells, and initiation of the immune response.

Cathepsin S, through proteolytic degradation of invariant chain to CLIP, provides a
20 fundamental step in generation of an immune response. It follows that inhibition of antigen presentation via prevention of invariant chain degradation by cathepsin S could provide a mechanism for immuno-regulation. Control of antigen-specific immune responses has long been desirable as a useful and safe therapy for autoimmune diseases. Such diseases include Crohn's disease and arthritis, as well as other T-cell-mediated
25 immune responses (C. Janeway and P. Travers, 1996, Immunobiology, The Immune System in Health and Disease, Chapter 12). Furthermore, cathepsin S, which has broad pH specificity, has been implicated in a variety of other diseases involving extracellular proteolysis, such as Alzheimer's disease (U. Muller-Ladner et al., 1996, Perspectives in Drug Discovery and Design, 6, 87) and atherosclerosis (G.K. Sukhova et al., 1998, J.
30 Clin. Invest., 102, 576).

A cathepsin S inhibitor has been found to block the rise in IgE titers and eosinophil infiltration in the lung in a mouse model of pulmonary hypersensitivity, suggesting that cathepsin S may be involved in asthma (R.J. Riese et al., J. Clin. Investigation, 1998, 101, 2351).

5

Another cysteine protease, cathepsin F has been found in macrophages and is also involved in antigen processing. It has been postulated that cathepsin F in stimulated lung macrophages and possibly other antigen presenting cells could play a role in airway inflammation (G.-P. Shi et al., J. Exp. Med., 2000, 191, 1177).

10

Cathepsin K, another cysteine protease has been found to be highly expressed in osteoclasts and to degrade bone collagen and other bone matrix proteins. Inhibitors of cathepsin K have been shown to inhibit bone resorption in mice. Therefore, cathepsin K may play a role in osteoclastic bone resorption and cathepsin K inhibitors may be useful in the treatment of diseases involving bone resorption such as osteoporosis (F. Lazner et al., Human Molecular Genetics, 1999, 8, 1839).

15

Cysteine proteases are characterized by having a cysteine residue at the active site which serves as a nucleophile. The active site also contains a histidine residue. The imidazole ring on the histidine serves as a base to generate a thiolate anion on the active site cysteine, increasing its nucleophilicity. When a substrate is recognized by the protease, the amide bond to be cleaved is directed to the active site, where the thiolate attacks the carbonyl carbon forming an acyl-enzyme intermediate and cleaving the amide bond, liberating an amine. Subsequently, water cleaves the acyl-enzyme species regenerating the enzyme and liberating the other cleavage product of the substrate, a carboxylic acid.

20

25

Inhibitors of cysteine proteases contain a functionality that can react reversibly or irreversibly with the active site cysteine. Examples of reactive functionalities that have been described (D. Rasnick, 1996, Perspectives in Drug Discovery and Design, 6, 47) on cysteine protease inhibitors include peptidyl diazomethanes, epoxides, monofluoroalkanes and acyloxymethanes, which irreversibly alkylate the cysteine thiol.

30

Other irreversible inhibitors include Michael acceptors such as peptidyl vinyl esters and other carboxylic acid derivatives (S. Liu et al., J. Med Chem., 1992, 35, 1067) and vinyl sulfones (J.T. Palmer et al., 1995, J. Med Chem., 38, 3193).

- 5 Reactive functionalities that form reversible complexes with the active site cysteine include peptidyl aldehydes (R.P. Hanzlik et al., 1991, Biochim. Biophys. Acta., 1073, 33), which are non-selective, inhibiting both cysteine and serine proteases as well as other nucleophiles. Peptidyl nitriles (R.P. Hanzlik et al., 1990, Biochim. Biophys. Acta., 1035, 62) are less reactive than aldehydes and therefore more selective for the more
- 10 nucleophilic cysteine proteases. Various reactive ketones have also been reported to be reversible inhibitors of cysteine proteases (D. Rasnick, 1996, *ibid*). In addition to reacting with the nucleophilic cysteine of the active site, reactive ketones may react with water, forming a hemiketal which may act as a transition state inhibitor.
- 15 Examples of cathepsin S inhibitors have been reported. J.L. Klaus et al. (WO 9640737) described reversible inhibitors of cysteine proteases including cathepsin S, containing an ethylene diamine. In US Patent No. 5,776,718 to Palmer et al. there is disclosed in it's broadest generic aspect a protease inhibitor comprising a targeting group linked through a two carbon atom chain to an electron withdrawing group (EWG). The compounds of the
- 20 present application are structurally distinct and thus excluded from the 5,776,718 patent with particular embodiments possessing unexpectedly greater activity than the closest compounds of the prior art. Other examples of cathepsin S inhibitors have been reported by E.T. Altmann et al, (WO 9924460, 1999) which describes dipeptide nitriles asserted to have activity as inhibitors of Cathepsins B, K, L and S. The WO publication does not
- 25 disclose any compounds possessing a succinate structure.

Additional peptidyl nitriles have been reported as protease inhibitors. For example, both nitriles and ketoheterocycles are described by B.A. Rowe et al. (US 5,714,471) as protease inhibitors useful in the treatment of neurodegenerative diseases. Peptidyl nitriles

30 are reported by B. Malcolm et al. (WO 9222570) as inhibitors of picornavirus protease.

B.J. Gour-Salin (Can. J. Chem., 1991, 69, 1288) and T.C. Liang (Arch. Biochim. Biophys., 1987, 252, 626) described peptidyl nitriles as inhibitors of papain

A reversible inhibitor presents a more attractive therapy than irreversible inhibitors.

- 5 Even compounds with high specificity for a particular protease can bind non-target enzymes. An irreversible compound could therefore permanently inactivate a non-target enzyme, increasing the likelihood of toxicity. Furthermore, any toxic effects resulting from inactivation of the target enzyme would be mitigated by reversible inhibitors, and could be easily remedied by modified or lower dosing. Finally, covalent modification of
- 10 an enzyme by an irreversible inhibitor could potentially generate an antibody response by acting as a hapten.

- In light of the above, there is a clear need for compounds which reversibly and selectively inhibit the cysteine proteases cathepsin S, K, F, L and B for indications in which these
- 15 proteases exacerbate disease.

SUMMARY OF THE INVENTION

- It is therefore an object of this invention to provide novel compounds according to the
- 20 formulas (I) and (Ia) as described herein which reversibly inhibit the cysteine proteases cathepsin S, K, F, L and cathepsin B. It is a further object of the invention to provide methods for treating diseases and pathological conditions exacerbated by these cysteine proteases such as, but not limited, to rheumatoid arthritis, asthma and osteoporosis. It is yet a further object of the invention to provide novel processes for preparation of the
- 25 above-mentioned novel compounds.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

30

The novel compounds of the invention are derivatives of succinamide and are therefore less peptidic in nature than many of the protease inhibitors described in the prior art.


These compounds, then, may exhibit certain advantages over known peptidic like compounds. These advantages may include, for example, increased bioavailability, increased stability, increased half-life and decreased clearance rates.

- 5 A proposed mechanism of action of the cysteine protease inhibitors of this invention is that the inhibitors contain a functionality that can react (reversibly or irreversibly) with the active site cysteine. The reactive functionality is attached to a peptide or peptide mimic that can be recognized and accommodated by the region of the protease surrounding the active site. The nature of both the reactive functionality and the remaining portion of the inhibitor determine the degree of selectivity and potency toward a particular protease.

Given the similarity of the active sites in cysteine proteases, it may be anticipated that a given class of inhibitors might have activity against more than one cysteine protease. It may also be expected that due to structural differences between individual cysteine proteases, different compounds of the invention may have different inhibitory potencies against different cysteine proteases. Thus some of the compounds of the invention may also be expected to be most effective in treating diseases mediated by cysteine proteases that they inhibit most potently. The activity of particular compounds disclosed herein against cysteine proteases cathepsin S, K, F, L and cathepsin B may be determined by the screens described in the section entitled "Assessment of Biological Properties."

Accordingly, in the first generic aspect of the invention, there is provided compounds of formula (I):

25



30 wherein:

A is -C(O)- or -CH(OR₈)-;

- 5 R₁ is alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl or amino wherein R₁ is optionally substituted by one or more R_a;

10 R_a is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, alkanoyl, aroyl, alkoxycarbonyl, aryloxycarbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, 15 arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_a may be further optionally substituted by one or more R_b;

20 R_b is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, 25 oxo, carboxy, cyano, nitro, amidino and guanidino;

R₂ is hydrogen, OR_i or lower alkyl;

30 R₃ is hydrogen or lower alkyl;

R₄ is hydrogen or lower alkyl;

35 R₅ is hydrogen, alkyl, cycloalkyl, aryl, heterocyclyl or heteroaryl, wherein R₅ is optionally substituted by one or more R_c;

40 R_c is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxycarbonyl, aryloxycarbonyl, alkanoyl, aroyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, 45 arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo,

carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_d ;

5 R_d is selected from the group consisting of alkyl, cycloalkyl, aryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

10 R_6 is hydrogen or lower alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S;

15 R_7 is hydrogen, alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S, cycloalkyl, aryl, heterocyclyl, aryl, heteroaryl or cyano, wherein R_7 is optionally substituted by one or more R_e ;

20 R_e is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, alkoxycarbonyl, aryloxycarbonyl, alkanoyloxy, aroyloxy, heteroarylalkoxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio, arylthio, arylalkylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_e may be further optionally substituted by one or more R_f ;

30 R_f is selected from the group consisting of alkyl, cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxycarbonyl, aryloxycarbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

45

or R6 and R7 together with the carbon they are attached form a 4 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, imidazolyl and pyridinyl, C1-5 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or disubstituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_g may be further optionally substituted by one or more R_h;

R_h is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, lower alkyl or lower alkoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranal and thiopyranal, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl and morpholinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl and pyridinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R8 is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl or aryl-alkyl;

Ri is hydrogen or lower alkyl;

X is O or S and

pharmaceutically acceptable salts, esters or tautomers thereof.

Preferred are compounds of the formula (I)

wherein:

5 A is as defined above;

R₁ is C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl and thiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl and amino wherein R₁ is optionally substituted by one or more R_a;

15 R_a is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by

alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, pyrazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl, aryl, C1-8 alkoxy, aryloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R₂ is hydrogen, OR_i or C1-5 alkyl;

R₃ is hydrogen or C1-5 alkyl;

R₄ is hydrogen or C1-5 alkyl;

R₅ is hydrogen, C1-8 alkyl, C3-7 cycloalkyl or aryl wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, alkanoyl, aroyl, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, ureido wherein either

nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxy carbonylamino, alkyl carbamoyloxy, aryl carbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl, aryl, arylalkyl, C1-8 alkoxy, aryloxy, arylalkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R_7 is hydrogen, C1-8 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S, C3-7 cycloalkyl, aryl or cyano, wherein R_7 is optionally substituted by one or more R_e ;

R_e is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, heteroarylC1-8alkoxy, C1-8 alkoxycarbonyl, aryloxy carbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl,

thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, arylC1-8 alkylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_e may be further optionally substituted by one or more R_f ,

R_f is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl,

5 isouquinoliny, quinazoliny, quinoxaliny, carbazolyl, phenothiaziny and
 phenoxaziny, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio,
 ureido wherein either nitrogen atom may be independently substituted by
 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidiny,
 10 piperidiny, morpholiny, thiomorpholiny, piperaziny and indoliny, or
 heteroaryl selected from the group consisting of furanyl, thienyl, pyrroly,
 oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl,
 oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridiny, pyridaziny,
 pyrimidiny, pyraziny, indolyl, isoindolyl, benzofuranyl, benzothienyl,
 15 benzimidazolyl, benzthiazolyl, puriny, quinoliny, isoquinoliny,
 quinazoliny, quinoxaliny, carbazolyl, phenothiaziny and phenoxaziny,
 alkoxycarbonylamino, aryloxy carbonylamino, alkyl carbamoyloxy,
 aryl carbamoyloxy, alkylsulfonylamino, arylsulfonylamino,
 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom
 20 may be independently mono or di-substituted by alkyl, aryl, heterocyclyl
 selected from the group consisting of pyrrolidiny, piperidiny,
 morpholiny, thiomorpholiny, piperaziny and indoliny, or heteroaryl
 selected from the group consisting of furanyl, thienyl, pyrroly, oxazolyl,
 thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl,
 triazolyl, tetrazolyl, thiadiazolyl, pyridiny, pyridaziny, pyrimidiny,
 pyraziny, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl,
 benzthiazolyl, puriny, quinoliny, isoquinoliny, quinazoliny,
 quinoxaliny, carbazolyl, phenothiaziny and phenoxaziny, halogen,
 25 hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

or R6 and R7 together with the carbon they are attached form a 4 to 7 membered
 carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally
 substituted with one or more R_g;

30 R_g is selected from the group consisting of alkyl, cycloalkyl, phenyl, heteroaryl selected
 from the group consisting of furanyl, thienyl and pyridiny, C1-5alkanoyl, aroyl, C1-
 5alkoxycarbonyl, aryloxy carbonyl, carbamoyl wherein the nitrogen atom may be
 independantly mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the
 35 group consisting of pyrrolidiny, piperidiny, morpholiny and piperaziny or heteroaryl
 selected from the group consisting of furanyl, thienyl, pyrroly, thiazolyl, imidazolyl,
 pyridiny, benzimidazolyl and quinoliny, C1-5 alkylthio wherein the sulfur atom may be
 oxidised to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidised to a
 40 sulfoxide or sulfone, C1-5 alkylaminosulfonyl, arylaminosulfonyl, halogen, hydroxy,
 oxo, carboxy and cyano,, R_g may be further optionally substituted by one or more R_h;

45 R_h is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl
 optionally substituted by one or more groups selected from halogen, C1-5 alkyl or
 C1-5 alkoxy, heterocyclyl selected from the group consisting of piperidiny,
 morpholiny and piperaziny, heteroaryl selected from the group consisting of

5 furanyl, thienyl pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo, carboxy and cyano;

10 R8 is hydrogen, alkyl, cycloalkyl-alkyl or arylalkyl;

15 Ri is hydrogen or C1-8 alkyl and

X is O or S.

20 More preferred are compounds of the formula (I) as described immediately above and wherein:

A is as defined above;

25 R1 is C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranal and thiopyranal, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl or amino; 30 wherein R1 is optionally substituted by one or more Ra;

35 Ra is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 40 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, 45

indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny and quinoxaliny, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny and quinoxaliny, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny and quinoxaliny, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R₂ is hydrogen, OR_i or C1-3 alkyl;

R₃ is hydrogen or C1-3 alkyl;

R₄ is hydrogen or C1-3 alkyl;

R₅ is hydrogen, C1-5 alkyl, C3-7 cycloalkyl or aryl wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny, isoquinoliny, quinazoliny and quinoxaliny, C1-5 alkoxy, aryloxy, C1-5 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl,

oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, amidino and guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, arylalkyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R₆ is hydrogen or C1-8 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

R₇ is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-7 cycloalkyl, aryl or cyano, wherein R₇ is optionally substituted by one or more R_e;

R_e is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, heteroarylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and

indolinyI or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyI, isoquinolinyI, quinazolinyI and quinoxalinyI, C1-5 alkanoylamino, aroylamino, C1-5 alkylthioarylthioarylC1-5 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyI or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyI, isoquinolinyI, quinazolinyI and quinoxalinyI, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyI or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyI, isoquinolinyI, quinazolinyI and quinoxalinyI, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_f,

R_f is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyI, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyI, isoquinolinyI, quinazolinyI and quinoxalinyI, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyI or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyI, isoquinolinyI, quinazolinyI and quinoxalinyI, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyI, or heteroaryl selected from the group consisting of furanyl,

thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy carbonylamino, aryloxy carbonylamino, C1-5 alkyl carbamoyloxy, aryl carbamoyloxy, C1-5 alkyl sulfonylamino, aryl sulfonylamino, C1-5 alkyl aminosulfonyl, aryl aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

or R₆ and R₇ together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl or a heterocyclic ring selected from the group consisting of pyrrolidinyl, piperidinyl, hexahydropyridazinyl, hexahydropyrimidinyl, piperazinyl, pyranyl, thiopyranyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide and imidazolidinyl-2,4-dione, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of alkyl, cycloalkyl, phenyl, heteroaryl selected from the group consisting of furanyl and thienyl, C1-3 alkanoyl, benzoyl, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, C1-3 alkylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, C1-5 alkyl aminosulfonyl, phenyl aminosulfonyl, halogen, hydroxy, carboxy and cyano, R_g may be further optionally substituted by one or more R_h;

R_h is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen, C1-3 alkyl and C1-3 alkoxy, piperidinyl, morpholinyl, piperazinyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, benzimidazolyl, quinolinyl, isoquinolinyl, C1-3 alkoxy, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo and cyano.

R₈ is hydrogen, C1-8 alkyl, C3-6 cycloalkyl-alkyl or aryl C1-3 alkyl; and

R_i is hydrogen or C1-5 alkyl.

Even more preferred are compounds of the formula (I) as described immediately above
5 and wherein:

A is as defined above;

10 R₁ is C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyranyl and thiopyranyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or amino wherein R₁ is optionally substituted by one or more R_a;

15 R_a is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group
25 consisting of pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl or aryl, C1-5
30 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from
35 the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy, cyano and nitro, R_a may be further
40 optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, halogen, hydroxy, oxo, carboxy and cyano;

45 R₂ is hydrogen, OR_i or methyl;

R3 is hydrogen or methyl;

R4 is hydrogen or methyl;

- 5 R5 is C1-5 alkyl, C3-7 cycloalkyl or phenyl wherein R5 is optionally substituted by one or more R_c;

10 R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen
15 atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido
20 wherein either nitrogen atom may be independently substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5
25 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further
30 optionally substituted by one or more R_d;

35

R_d is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-5alkyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy and cyano;

40

R6 is hydrogen or C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

45

R₇ is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-7 cycloalkyl, phenyl, naphthyl or cyano, wherein R₇ is optionally substituted by one or more R_c;

5 R_c is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl
10 selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, heteroarylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be
15 independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, arylC1-5 alkylthio, ureido wherein either
20 nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5
25 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected
30 from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_f;

35 R_f is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen, methyl and methoxy, naphthyl optionally substituted by one or more groups selected from halogen, methyl and methoxy, heterocyclyl
40 selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl,
45 C1-5 alkoxy, aryloxy, arylC1-5alkoxy, C1-5 alkoxycarbonyl,

aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano;

or R6 and R7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl or a heterocyclic ring selected from the group consisting of piperidinyl, hexahydropyridazinyl, hexahydropyrimidinyl, piperazinyl, pyranyl and thiopyranyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, C1-3 alkanoyl, benzoyl, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, carboxy and cyano, R_g may be further optionally substituted by one or more R_h;

R_h is selected from the group consisting of methyl, phenyl optionally substituted by one or more groups selected from halogen, methyl and methoxy, piperidinyl, morpholinyl, piperazinyl, pyridinyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo and cyano;

R₈ is hydrogen, C1-5 alkyl, C5-6 cycloalkyl-C1-3-alkyl or benzyl;

R_i is hydrogen or methyl; and

5 X is O.

Yet even more preferred are compounds of the formula (I) as described immediately above and wherein:

10

A is as defined above;

15 R₁ is C1-5 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl, piperazinyl, pyranyl and thiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or amino, wherein R₁ is optionally substituted by one or more R_a;

20 R_a is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, 25 quinolinyl and isoquinolinyl, C1-3 alkoxy, phenoxy, C1-3 alkoxycarbonyl, aryloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 30 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5alkyl, phenyl or naphthyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 35 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, 40 halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

45 R_b is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, aryl, C1-3 alkoxy, phenoxy, halogen, hydroxy, oxo, carboxy and cyano;

R2 is hydrogen or ORi;

R3 is hydrogen;

5 R4 is hydrogen;

R5 is C1-5 alkyl, C3-6 cycloalkyl or phenyl, wherein R5 is optionally substituted by one or more R_c;

10

15

20

25

30

R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, phenoxy, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio, phenylthio, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

35

R_d is selected from the group consisting of C1-5 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, arylC1-3alkyl, C1-5 alkoxy, phenoxy, arylC1-3alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy and cyano;

40

R6 is hydrogen or C1-3alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

R7 is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-6 cycloalkyl, phenyl or cyano, wherein R7 is optionally substituted by one or more R_e;

45

R_e is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl,

piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, heteroarylC1-3alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio, arylC1-3alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-5 alkoxycarbonylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, and indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen and methyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl and indolyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or aryl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, halogen, hydroxy, oxo, carboxy and cyano;

or

R₆ and R₇ together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl or heterocyclic ring selected from the group consisting of piperidinyl, morpholinyl and thiomorpholinyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, C1-3 alkanoyl, C1-3 alkoxycarbonyl and carbamoyl wherein the nitrogen atom may be

independantly mono or disubstituted by C1-5 alkyl; R_g may be further optionally substituted by one or more R_h ;

5 R_h is phenyl or amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl;

R_8 is hydrogen or methyl;

10 R_i is hydrogen or methyl and
and X is O.

15 Still yet even more preferred are compounds of the formula (I) as described immediately above and wherein:

A is as defined above;

20 R_1 is C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl, piperazinyl, pyranyl and thiopyranyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl or amino, wherein R_1 is optionally substituted by one or more R_a ;

25 R_a is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-3 alkoxy, C1-3
30 alkoxy carbonyl, aryloxy carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl or heteroaryl selected from the group consisting of pyrrolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl and benzthiazolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio, arylthio, ureido
35 wherein either nitrogen atom may be independently substituted by C1-3alkyl or phenyl, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of pyrrolyl,
40 imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b ;

45 R_b is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, C1-3 alkoxy, halogen and hydroxy;

R2 is hydrogen;

R3 and R4 are as defined immediately above;

- 5 R5 is C1-5 alkyl, C5-6 cycloalkyl or phenyl wherein R5 is optionally substituted by one or more R_c;

10 R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, 4-piperidinyl, 4-morpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-5 alkoxy, phenoxy, aroyl, C1-5 alkoxycarbonyl, aryloxy carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, C1-5 alkanoylamino, C1-3 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or phenyl, C1-5 alkoxycarbonylamino, 15 C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

20 R_d is selected from the group consisting of C1-5 alkyl, C5-6 cycloalkyl, phenyl, benzyl, C1-5 alkoxy, phenoxy, benzyloxy, aroyl, halogen, hydroxy, oxo, carboxy and cyano;

- 25 R6 is hydrogen or C1-3alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

30 R7 is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, phenyl or cyano, wherein R7 is optionally substituted by one or more R_e;

35 R_e is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, benzyloxy, pyridylC1-3alkoxy, thienylC1-3alkoxy, furanylC1-3alkoxy, C1-3 alkoxycarbonyl, phenoxyoxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, methylthio, benzylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-3 alkoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, halogen, hydroxy, 40 oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

45 R_f is selected from the group consisting of C1-3 alkyl, phenyl optionally substituted by one or more groups selected from the group consisting of

halogen and methyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl and pyridinyl, C1-3 alkoxy, aryloxy, benzyloxy, C1-5 alkoxy carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-5 alkoxy carbonylamino, C1-5 alkyl carbamoyloxy, aryl carbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano;

or R6 and R7 together with the carbon they are attached form cyclopropyl or a heterocyclic ring selected from the group consisting of piperidinyl, pyranal and thiopyranal, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of methyl, benzyl, acetyl, benzoyl, benzyloxy carbonyl and ethoxy carbonyl;

R₈ is hydrogen and

X is O.

Even much more preferred are compounds of the formula (I) as described immediately above and wherein:

A is as described above;

R₁ is C3-6 cycloalkyl, phenyl, naphthyl, piperidinyl, morpholinyl, piperazinyl, pyranal, thiopyranal, furanyl, thienyl, pyrrolyl or amino, wherein R₁ is optionally substituted by one or more R_a;

R_a is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or phenyl, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-3alkyl, C1-3alkoxy, halogen and hydroxy;

5 R₂, R₃ and R₄ are as defined immediately above;

R₅ is C1-5 alkyl, C5-6 cycloalkyl or phenyl, wherein R₅ is optionally substituted by one or more groups of the formula R_c;

10 R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, 4-piperidinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, C1-5 alkoxy carbonyl, C1-5 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, C1-3 alkylthio, C1-3 alkoxy carbonylamino, 15 C1-3 alkylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

20 R_d is selected from the group consisting of C1-3 alkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, benzyloxy, benzoyl, halogen, hydroxy, oxo, carboxy and cyano;

wherein the configuration at the stereocenter defined by R₄ and R₅ and the carbon they are attached to is defined as L;

25 R₆ is hydrogen or C1-2 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S;

30 R₇ is C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S; phenyl or cyano wherein R₇ is optionally substituted by one or more groups of the formula R_e;

35 R_e is selected from the group consisting of methyl, C3-6 cycloalkyl, phenyl, naphthyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, benzyloxy, pyridyl, C1-3alkoxy, thienyl, C1-3alkoxy, furanyl, C1-3alkoxy, C1-5 alkanoylamino, aroylamino, methylthio, benzylthio, C1-3 alkoxy carbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f.

40 R_f is selected from the group consisting of C1-3 alkyl, phenyl optionally substituted by one or more groups selected from halogen and methyl, C1-3 alkoxy, aryloxy, benzyloxy, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, amino wherein the nitrogen 45

atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano;

or R6 and R7 together with the carbon they are attached form cyclopropyl or a heterocyclic ring selected from the group consisting of piperidinyl and pyranyl, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g.

R_g is methyl;

R_h is as described immediately above; and

X is O.

Yet even much more preferred are compounds of the formula (I) as described immediately above and wherein:

A is as described above;

R1 is cyclopropyl, cyclohexyl, phenyl, naphthyl, piperidinyl, morpholinyl, piperazinyl, pyranyl, thiopyranyl or amino wherein R1 is optionally substituted by one or more R_a;

R_a is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-3 alkoxy, halogen and hydroxy,

R2, R3 and R4 are as described immediately above;

R5 is C1-5 alkyl or C5-6 cycloalkyl wherein R5 is optionally substituted by one or more R_c;

R_c is selected from the group consisting of methyl, C3-6 cycloalkyl, phenyl, naphthyl, thienyl, imidazolyl, pyridinyl, indolyl, C1-4 alkoxy, C1-5 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of methyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, halogen and hydroxy;

R_6 is as described immediately above;

R_7 is C1-5 alkyl or phenyl, wherein R_7 is optionally substituted by one or more R_e ;

R_e is selected from the group consisting of C3-6 cycloalkyl, phenyl, naphthyl, thienyl, imidazolyl, pyridinyl, indolyl, methoxy, benzyloxy, C1-3 alkanoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, benzylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, methoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f ;

R_f is selected from the group consisting of methyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, methoxy, phenoxy, benzyloxy, methoxycarbonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy and carboxy;

or R_6 and R_7 together with the carbon they are attached form a cyclopropyl ring; and

R_g , R_h and X are as described immediately above.

Penultimately preferred are compounds of the formula (I) as described immediately above and wherein:

A is as described above;

R_1 is cyclopropyl, cyclohexyl, phenyl, naphthyl, piperidinyl, morpholinyl, piperazinyl, pyranyl or thiopyranyl, wherein R_1 is optionally substituted by one or more R_a ;

R_a is selected from the group consisting of pyrrolyl, imidazolyl, indolyl, benzimidazolyl, methoxy, methoxycarbonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy and carboxy, R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of methoxy, halogen and hydroxy;

R_2, R_3, R_4 and R_5 are as described immediately above;

R_c is selected from the group consisting of methyl, C3-6 cycloalkyl, phenyl, naphthyl, C1-4 alkoxy, C1-3 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

5 R_d is selected from the group consisting of methyl, phenyl, methoxy, halogen and hydroxy;

R₆ is hydrogen;

10 R₇ is as described immediately above;

 R_e is selected from the group consisting of C5-6 cycloalkyl, phenyl, naphthyl, thienyl, indolyl, methoxy, benzyloxy, methylthio, benzylthio, methoxycarbonylamino, halogen, hydroxy, carboxy and cyano, R_e may be further
15 optionally substituted by one or more R_f;

 R_f is selected from the group consisting of methyl, phenyl optionally substituted by halogen, methoxy, phenoxy, benzyloxy, methoxycarbonyl, halogen, hydroxy and carboxy; and
20

R_g, R_h and X are as described immediately above.

25

Ultimately preferred are compounds of the formula(I) wherein:

A is as described above;

30

R₁ is 4-morpholinyl, 4-pyranyl or phenyl;

R₂, R₃, R₄, R_a and R_b are as described immediately above;

35

R₅ is C1-5 alkyl wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C3-6 cycloalkyl, phenyl and 2-naphthyl, R_c may be further optionally substituted by one or more R_d;

40

R_d is selected from the group consisting of methyl and halogen,

R₆ and R₇ are as described immediately above;


45

R_e is selected from the group consisting of C5-6 cycloalkyl, phenyl, naphthyl, indolyl, benzyloxy, methylthio, benzylthio, halogen and carboxy, R_e may be further optionally substituted by one or more R_f; and

R_f is selected from the group consisting of methyl, methoxy, methoxycarbonyl, halogen and hydroxy; and

5 R_g , R_h and X are as described immediately above.

In a second generic aspect of the invention, there are provided compounds of formula
10 (Ia):



wherein:

A is $-C(O)-$ or $-CH(OR_8)-$;

R_1 is alkyl, alkoxy, cycloalkyl, aryl, aryloxy, benzyloxy, heterocyclyl, heteroaryl or amino wherein R_1 is optionally substituted by one or more R_a ;

R_a is selected from the group consisting of a bond, alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, alkanoyl, aroyl, alkoxycarbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_a may be further optionally substituted by one or more R_b ;

5 R_8 is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, acyl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R2 is hydrogen, OR_j or lower alkyl;

10 R3 is hydrogen or lower alkyl;

R4 is hydrogen or lower alkyl;

15 R5 is hydrogen, alkyl, cycloalkyl, aryl, heterocyclyl or heteroaryl, wherein R5 is optionally substituted by one or more R_c ;

20 R_c is selected from the group consisting of a bond, alkyl, cycloalkyl, bicycloalkyl, aryl, indenyl, indanyl, dihydronaphthyl, tetrahydronaphthyl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxycarbonyl, aryloxy carbonyl, alkanoyl, aroyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_d ;

35 R_d is selected from the group consisting of alkyl, cycloalkyl, aryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

40 or R4 and R5 together with the carbon they are attached form a 3 to 7 membered carbocyclic ring optionally substituted with one or more R_d ;

R6 is hydrogen or lower alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S;

45

R7 is hydrogen, alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S, cycloalkyl, aryl, heterocyclyl, heteroaryl or cyano, wherein R7 is optionally substituted by one or more R_e;

5

R_e is selected from the group consisting of a bond alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, alkoxy carbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, heteroarylalkoxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylalkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxy carbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_e may be further optionally substituted by one or more R_f;

10

R_f is selected from the group consisting of alkyl, cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxy carbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxy carbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

25

30

35

40

or R6 and R7 together with the carbon they are attached form a 3 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

45

R_g is selected from the group consisting of a bond, alkyl, cycloalkyl, bicycloalkyl, benzyl, aryl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, tetrahydrothiopyranyl, indolinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrimidinyl, pyrrolyl, imidazolyl and pyridinyl, C1-5 alkanoyl, aroyl, C1-5 alkoxy carbonyl, aryloxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or disubstituted by alkyl, cycloalkyl, bicycloalkyl, aryl, aroyl, benzoyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_g may be further optionally substituted by one or more R_h ;

R_h is selected from the group consisting of a bond, C1-5 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, lower alkyl or lower alkoxy, benzyl, benzyloxy, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl and pyridinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_h may be further optionally substituted by one or more R_i ;

R_i is alkyl, hydroxy, oxo and halogen;

R_j is hydrogen, alkyl, cycloalkyl;

R_8 is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl or arylalkyl;

X is O or S;

and

pharmaceutically acceptable salts, esters or tautomers thereof.

5

Preferred are compounds of the formula (Ia)

10 wherein:

15 R₁ is C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, 2-oxa-5-aza-bicyclo[2,2,1]heptanyl, piperazinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, 20 isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl and amino wherein R₁ is optionally substituted by one or more R_a;

25 R_a is selected from the group consisting of a bond, C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, 30 benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and 35 indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, 40 C1-8 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, 45 heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl,

morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, pyrazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino, guanidino; R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl, aryl, heteroaryl, C1-8 alkoxy, aryloxy, alkanoyl, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R_2 is hydrogen, OR_j or C1-3 alkyl;

R_3 is hydrogen or C1-5 alkyl;

R_4 is hydrogen or C1-5 alkyl;

R_5 is hydrogen, C1-8 alkyl, C3-7 cycloalkyl, aryl wherein R_5 is optionally substituted by one or more R_c ;

R_c is selected from the group consisting of a bond, C1-8 alkyl, C3-7 cycloalkyl, bicycloalkyl, aryl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl,

morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, alkanoyl, aroyl, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, C1-5 aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl, aryl, arylalkyl, C1-8 alkoxy, aryloxy, arylalkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R7 is hydrogen, C1-8 alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-7 cycloalkyl, aryl, heteroaryl or cyano, wherein R7 is optionally substituted by one or more R_e;

5 R_e is selected from the group consisting of a bond, C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, 10 pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, heteroarylC1-8alkoxy wherein the heteroaryl is as hereinabove described in this paragraph, C1-8 alkoxycarbonyl, aryloxycarbonyl, 15 C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, 20 isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylC1-8 alkylthio, ureido wherein either 25 nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, 30 alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, 40 oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, 45 halogen, hydroxy, oxo, carboxy, cyano, nitro,

amidino and guanidino, R_c may be further optionally substituted by one or more R_f

5 R_f is selected from the group consisting C1-8 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from the group consisting halogen, methyl and methoxy, heterocyclyl selected from the group consisting pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of
 10 furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, C1-8 alkoxy carbonyl, aryloxy carbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-
 15 substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8
 20 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxy carbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein
 30 the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxy carbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein
 35 the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl,
 40
 45

benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino, and guanidino;

or R₆ and R₇ together with the carbon they are attached form a 3 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of a bond, alkyl, cycloalkyl, bicycloalkyl, phenyl, naphthyl, benzyl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, tetrahydrothiopyranyl, indolinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrimidinyl and pyridinyl, C1-5alkanoyl, aroyl, C1-5alkoxycarbonyl, aryloxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, benzyl, benzoyl or naphthoyl, C1-5 alkylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, C1-5 alkylaminosulfonyl, arylaminosulfonyl, halogen, hydroxy, oxo, carboxy and cyano, R_g may be further optionally substituted by one or more R_h;

R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, C1-5 alkyl or C1-5 alkoxy, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl, or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo, carboxy and cyano; and

R₈ is hydrogen, alkyl, cycloalkyl-alkyl or arylalkyl;

R_i is C1-8 alkyl, hydroxy, oxo and halogen

R_j is hydrogen or alkyl and

X is O.

5

More preferred are compounds of the formula (I) as described immediately above and wherein:

10

A is $-\text{C}(\text{O})-$;

15

R₁ is C1-5 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, 2-oxa-5-aza-bicyclo[2,2,1]heptanyl, piperazinyl, tetrahydropyranyl and tetrahydrothiopyranyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or amino, wherein R₁ is optionally substituted by one or more R_a;

20

25

30

35

40

R_a is selected from the group consisting of a bond, C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-3 alkoxy, phenoxy, C1-3 alkoxycarbonyl, aryloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, phenyl or naphthyl; C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

45

R_b is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, aryl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl,

benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl,
C1-3 alkoxy, phenoxy, C1-8 alkanoyl, amino wherein the nitrogen atom
may be independently mono or di-substituted by C1-5alkyl, C3-6
cycloalkyl, phenyl or naphthyl, halogen, hydroxy, oxo, carboxy and
cyano;

R2 is hydrogen;

R3 is hydrogen;

R4 is hydrogen;

R5 is C1-5 alkyl, C3-6 cycloalkyl or phenyl, wherein R5 is optionally substituted by one
or more R_c;

R_c is selected from the group consisting of a bond, C1-3 alkyl, C3-6 cycloalkyl,
phenyl, naphthyl, indenyl, indanyl, heterocyclyl selected from the group
consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from
the group consisting of furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl,
quinolinyl and isoquinolinyl, C1-5 alkoxy, phenoxy, aroyl, C1-5 alkoxycarbonyl,
aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen
atom may be independently mono or di-substituted by C1-5 alkyl, phenyl,
naphthyl, heterocyclyl selected from the group consisting of piperidinyl,
morpholinyl and piperazinyl or heteroaryl selected from the group consisting of
furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl,
C1-5 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be
oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
independently substituted by C1-3 alkyl, phenyl or heteroaryl selected the group
consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl
and indolyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy,
arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein
the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl,
phenyl, heterocyclyl selected from the group consisting of piperidinyl,
morpholinyl and piperazinyl or heteroaryl selected from the group consisting of
furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl,
halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally
substituted by one or more R_d;

R_d is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl,
phenyl, naphthyl, arylC1-3alkyl, C1-5 alkoxy, phenoxy, arylC1-3alkoxy,
aroyl, amino, halogen, hydroxy, oxo, carboxy and cyano;

R₆ is hydrogen or C1-3alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S;

5 R₇ is hydrogen, C1-5 alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-6 cycloalkyl, phenyl, naphthyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or cyano, wherein R₇ is optionally substituted by one or more R_e;

10 R_e is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, 15 indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, heteroarylC1-3alkoxy wherein the heteroaryl is as hereinabove described in this paragraph, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 20 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylC1-3alkylthio wherein the sulfur atom may 25 be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-5 alkoxycarbonylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and 30 piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl and pyridinyl, indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

35 R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, 40 imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl and indolyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may 45 be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom

may be independently substituted by C1-5 alkyl or aryl, C1-5 alkoxy carbonylamino, aryloxy carbonylamino, C1-5 alkyl carbamoyloxy, aryl carbamoyloxy, C1-5 alkyl sulfonylamino, aryl sulfonylamino, C1-5 alkyl aminosulfonyl, aryl aminosulfonyl, amino wherein the nitrogen atom
5 may be independently mono or di-substituted by C1-5 alkyl or aryl, halogen, hydroxy, oxo, carboxy and cyano;

or

- 10 R₆ and R₇ together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or heterocyclic ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, tetrahydropyrany and tetrahydrothiopyrany, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

- 15 R_g is selected from the group consisting of a bond, C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidiny, piperidiny, tetrahydropyrany, tetrahydrothiopyrany, piperaziny and morpholiny, heteroaryl
20 selected from the group consisting of thienyl, pyrimidiny and pyridiny, C1-5 alkanoyl, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, benzyl, benzoyl or naphthoyl, R_g may be
25 further optionally substituted by one or more R_h; and

- R_h is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, phenyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of piperidiny, indoliny, tetrahydropyrany and tetrahydrothiopyrany, heteroaryl selected from
30 the group consisting of thienyl, pyrrolyl, pyridiny, pyrimidiny, indolyl, benzimidazolyl, quinoliny and isoquinoliny, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, oxo, cyano and trifluoromethyl.

- 35 Even more preferred are compounds of the formula (Ia) as desribed immediately above and wherein:

- 40 R₁ is C3-6 cycloalkyl, phenyl, naphthyl, pyrrolidiny, piperidiny, morpholiny, thiomorpholiny, piperaziny, tetrahydropyrany, furany, thienyl, pyrrolyl or amino, wherein R₁ is optionally substituted by one or more R_a;

- R_a is selected from the group consisting of a bond, C1-3 alkyl, C5-6 cycloalkyl, phenyl, furany, thienyl, pyrrolyl, imidazolyl, indolyl, benzofurany, benzothienyl,
45 benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl;

C1-5alkoxycarbonylamino, C1-5 alkanoylamino, aroylamino, ureido wherein either nitrogen atom may be independently substituted by C1-3alkyl or phenyl; C1-5 alkylsulfonfylamino, arylsulfonfylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-3alkyl, C1-3alkoxy, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3alkyl or phenyl halogen, oxo and hydroxy;

R₅ is C1-5 alkyl, C3-6 cycloalkyl or phenyl, wherein R₅ is optionally substituted by one or more groups of the formula R_c;

R_c is selected from the group consisting of a bond, C1-3 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, 4-piperidiny, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, C1-5 alkoxycarbonyl, C1-5 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-3 alkoxycarbonylamino, C1-3 alkylsulfonfylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of C1-3 alkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, benzyloxy, benzoyl, halogen, hydroxy, oxo, carboxy and cyano;

wherein the configuration at the stereocenter defined by R₄ and R₅ and the carbon they are attached to is defined as L;

R₆ is hydrogen or C1-2alkyl one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S;

R₇ is C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S; phenyl, pyridinyl or cyano wherein R₇ is optionally substituted by one or more groups of the formula R_e;

R_e is selected from the group consisting of a bond, methyl, C3-6 cycloalkyl, phenyl, naphthyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrrolidinyl, piperidinyl, indolyl, C1-3 alkoxy, benzyloxy, pyridinylC1-3alkoxy, thienylC1-3alkoxy, furanylC1-3alkoxy, C1-5 alkanoylamino, aroylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, benzylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-3

alkoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f .

5 R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, C1-3 alkoxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-
10 substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano;

15 R_6 and R_7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclopentyl, cyclohexyl and cycloheptyl or heterocyclic ring selected from the group consisting of pyrrolidinyl, piperidinyl and tetrahydropyranyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g ;

20 R_g is selected from the group consisting of a bond C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, dihydronaphthyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from the group consisting of piperidinyl, tetrahydropyranyl, and tetrahydrothiopyranyl, C1-5 alkanoyl, C1-3 alkoxycarbonyl, carbamoyl
25 wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, benzyl, benzoyl or naphthoyl, R_g may be further optionally substituted by one or more R_h ; and

30 R_h is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, phenyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of piperidinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of thienyl, pyridinyl, indolyl, pyrrolyl and benzimidazolyl,
35 amino wherein the nitrogen atom may be independently mono or di-substituted by methyl, halogen, hydroxy, oxo, cyano and trifluoromethyl.

40 Yet even more preferred are compounds of the formula (Ia) as described immediately above and wherein:

45 R_1 is cyclohexyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, or amino wherein R_1 is optionally substituted by one or more R_a ;

R_a is selected from the group consisting of a bond, C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkoxycarbonylamino, C1-5 alkanoylamino, aroylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-3 alkoxy, halogen and hydroxy,

R_5 is C1-5 alkyl wherein R_5 is optionally substituted by one or more R_c ;

R_c is selected from the group consisting of a bond, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, indanyl, thienyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, C1-5 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of C1-3 alkyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, halogen and hydroxy;

R_7 is C1-5 alkyl or phenyl, wherein R_7 is optionally substituted by one or more R_e ;

R_e is selected from the group consisting of a bond, C3-6 cycloalkyl, phenyl, naphthyl, thienyl, imidazolyl, pyrrolidinyl, piperidinyl, pyridinyl, indolyl, C1-5 alkoxy, benzyloxy, C1-5 alkanoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, benzylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, methoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, halogen, hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f ;

R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, methoxy, phenoxy, benzyloxy, methoxycarbonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy and carboxy;

R_6 and R_7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl and cyclohexyl or heterocyclic ring selected from the group consisting of pyrrolidinyl and piperidinyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g ;

5 R_g is selected from the group consisting of a bond, C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from the group consisting of tetrahydropyranyl and tetrahydrothiopyranyl, acetyl, methoxycarbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-3 alkyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, benzyl or benzoyl, R_g may be further optionally substituted by one or more R_h ; and

10 R_h is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, phenyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of piperidinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of thienyl, pyridinyl, indolyl and benzimidazolyl, amino, methylamino, dimethylamino, halogen, hydroxy, oxo, cyano and trifluoromethyl.

15 Still yet even more preferred are compounds of the formula (I) as described immediately above and wherein:

20 R_1 is phenyl, naphthyl, pyrrolidinyl, thiomorpholinyl or morpholinyl, wherein R_1 is optionally substituted by one or more R_a ;

25 R_a is selected from the group consisting of a bond, C1-3 alkyl, pyrrolyl, imidazolyl, indolyl, benzimidazolyl, methoxy, methoxycarbonyl, t-butoxycarbonylamino, C1-3 alkylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, cyano and carboxy, R_a may be further optionally substituted by one or more R_b ;

30 R_b is selected from the group consisting of methoxy, halogen and hydroxy;

35 R_c is selected from the group consisting of a bond, methyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, tetrahydronaphthyl, indanyl, C1-4 alkoxy, C1-3 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d ;

40 R_d is selected from the group consisting of C1-3 alkyl, phenyl, methoxy, halogen and hydroxy;

R_6 is hydrogen;

45 R_e is selected from the group consisting of a bond, C5-6 cycloalkyl, phenyl, C1-5 alkoxy, C1-5 alkylamino, pyrrolidinyl, piperidinyl, naphthyl, thienyl, indolyl, methoxy, benzyloxy, methylthio, benzylthio, methoxycarbonylamino, halogen,

hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f , and

5 R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by halogen, methoxy, phenoxy, benzyloxy, methoxycarbonyl, halogen, hydroxy and carboxy.

10 R_6 and R_7 together with the carbon they are attached form a cyclopropyl ring or heterocyclic ring selected from the group consisting of pyrrolidinyl and piperidinyl, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g ;

15 R_g is selected from the group consisting of a bond, C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, α -tetrahydronaphthyl, α -indanyl, tetrahydropyranyl, tetrahydrothiopyranyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, benzyl or benzoyl, R_g may be further optionally substituted by one or more R_h ; and

20 R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, benzyl, benzyloxy, tetrahydropyranyl, heteroaryl selected from the group consisting of thienyl, pyridinyl, indolyl, pyrrolyl and benzimidazolyl, halogen, hydroxy, oxo, cyano or trifluoromethyl, may be further substituted by R_i ; and

R_i is C1-8 alkyl and halogen.

25

Even much more preferred are compounds of the formula (Ia) as described immediately above and wherein:

30

R_1 is phenyl, morpholinyl, thiomorpholinyl or pyrrolidinyl, R_1 is optionally substituted by one or more R_a ;

35 R_a is selected from the group consisting of C1-3 alkyl, methoxy, C1-3 alkylsulfonylamino, halogen, hydroxy, cyano and trifluoromethyl;

R_5 is C1-3 alkyl wherein R_5 is optionally substituted by one or more R_c ;

40 R_c is selected from the group consisting of C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, β -tetrahydronaphthyl or β -indanyl, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of methyl, methoxy and halogen;

45 R_7 is C1-5 alkyl wherein R_7 is optionally substituted by one or more R_e ;

R_e is selected from the group consisting of phenyl, naphthyl, C1-5 alkoxy, C1-5 alkylamino, pyrrolidinyl and piperidinyl, R_e may be further optionally substituted by one or more R_f ; and

5 R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted with halogen, halogen and hydroxy.

or R_6 and R_7 together with the carbon they are attached form a cyclopropyl ring or heterocyclic ring selected from the group consisting of piperidinyl and pyrrolidinyl, the
10 cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g ; and

R_g is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, α -tetrahydronaphthyl, α -indanyl, tetrahydropyranyl, tetrahydrothiopyranyl and amino group wherein the nitrogen atom may be
15 independantly mono or disubstituted by C1-5 alkyl or C3-6 cycloalkyl; R_g may be further optionally substituted by one or more R_h ; and

R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, benzyl, benzyloxy, thienyl, pyrrolyl, indolyl, pyridinyl, halogen, hydroxy, cyano and
20 trifluoromethyl; R_h may be further optionally substituted by one or more R_i ; and

R_i is C1-5 alkyl and halogen.

25 Yet even much more preferred are compounds of the formula (Ia) as described immediately above and wherein:

R_1 is 4-morpholinyl or pyrrolidinyl, wherein R_1 is optionally substituted by one or more
30 R_a ;

R_a is C1-3 alkylsulfonylamino;

R_5 is C1-3 alkyl wherein R_5 is optionally substituted by one or more R_c ;

35 R_c is selected from the group consisting of C5-6 cycloalkyl, C7-8 bicycloalkyl, β -tetrahydronaphthyl and β -indanyl, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of methyl, methoxy and halogen;

40 R_7 is C1-5 alkyl wherein R_7 is optionally substituted by one or more R_e .

R_e is selected from the group consisting of C1-5 alkoxy, C1-5 alkylamino, pyrrolidinyl and piperidinyl, R_e may be further optionally substituted by one or more R_f; and

R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted with halogen, halogen and hydroxy.

or R₆ and R₇ together with the carbon they are attached form heterocyclic ring selected from the group consisting of piperidinyl and pyrrolidinyl, the heterocyclic ring being optionally substituted with one or more R_g; and

R_g is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, α -tetrahydronaphthyl, α -indanyl, tetrahydropyranyl and tetrahydrothiopyranyl; R_g may be further optionally substituted by one or more R_h; and

R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, pyridinyl, indolyl, thienyl, halogen, hydroxy, cyano and trifluoromethyl; R_h may be further optionally substituted by one or more R_i;

R_i is selected from the group consisting of C1-3 alkyl and halogen.

Ultimately preferred are compounds of the formula (Ia) as described immediately above and wherein:

R₁ is 4-morpholinyl;

R₅ is C1-3 alkyl wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of cyclohexyl, β -tetrahydronaphthyl and β -indanyl, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of methyl and halogen;

R₇ is C1-5 alkyl wherein R₇ is optionally substituted by one or more R_e;

R_e is selected from the group consisting of C1-2alkoxy and C1-2 alkylamino, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of C1-3 alkyl, phenyl, halogen and hydroxy;

5 or R6 and R7 together with the carbon they are attached form a heterocyclic ring selected from the group consisting of piperidinyl and pyrrolidinyl, the heterocyclic ring being optionally substituted with one or more R_g; and

10 R_g is selected from the group consisting of C1-5 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl and α -tetrahydronaphthyl; R_g may be further optionally substituted by one or more R_h; and

R_h is C1-5 alkyl, C3-7 cycloalkyl, thienyl, pyridinyl, indolyl, halogen, hydroxy and trifluoromethyl; R_h may be further optionally substituted by one or more R_i;

15 R_i is methyl.

The activity of particular compounds disclosed herein against cathepsin K may be determined without undue experimentation by one of ordinary skill in the art in view of
20 the art, the guidance provided throughout this specification and by the art recognized methods referred to in the section entitled "Assessment of Biological Properties."

The following subgeneric aspect of the compounds of the formula (Ia) is postulated to possess Cathepsin K activity:

25

The broadest embodiment of the formula (Ia) as described hereinabove and wherein

A is -C(O)-;

30 R1 is aryl or aryloxy wherein R1 is optionally substituted by one or more R_a;

R2 is hydrogen;

R3 is hydrogen;

35 R4 is hydrogen or lower alkyl;

R5 is alkyl optionally substituted by one or more R_c;

5 or R4 and R5 together with the carbon they are attached form a 3 to 7 membered carbocyclic ring optionally substituted with one or more R_d;

R6 is hydrogen or lower alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S;

10 R7 is hydrogen, alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S, cycloalkyl, aryl, heterocyclyl, heteroaryl or cyano, wherein R7 is optionally substituted by one or more R_e;

15 R6 and R7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or heterocyclic ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, tetrahydropyrany and tetrahydrothiopyrany, the carbocyclic or
20 heterocyclic ring being optionally substituted with one or more R_g;

Preferred cathepsin K inhibitors are those as described immediately above and wherein:

25 R1 is naphthyl, benzyloxy or phenoxy wherein R1 is optionally substituted by one or more R_a;

R4 is hydrogen or lower alkyl;

30 R5 is lower alkyl optionally substituted by one or more R_c;

or R4 and R5 together with the carbon they are attached form a 5 to 6 membered carbocyclic ring optionally substituted with one or more R_d;

35 R6 is hydrogen or lower alkyl wherein one or more carbon atoms are optionally replaced by N;

40 R7 is lower alkyl wherein one or more carbon atoms are optionally replaced by heteroatoms selected from the group consisting of N and O, R7 is optionally substituted by one or more R_e;

R6 and R7 together with the carbon they are attached form a heterocyclic ring selected from the group consisting of pyrrolidinyl and piperidinyl each being optionally substituted with one or more R_g .

5

Most preferred cathepsin K inhibitors are those as described immediately above and wherein:

10 R_4 is hydrogen or lower alkyl;

R_5 is C1-4 alkyl optionally substituted by one or more R_c ;

15 or R_4 and R_5 together with the carbon they are attached form a cyclohexyl ring optionally substituted with one or more R_d ;

R_7 is lower alkyl wherein one or more carbon atoms are optionally replaced by heteroatoms selected from the group consisting of N and O or benzyloxy;

20

R_6 and R_7 together with the carbon they are attached form pyrrolidinyl optionally substituted with one or more R_g .

25

The following are representative compounds of the invention:

4-Methyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (2-benzyloxy-1-cyano-ethyl)-amide. MS: $m/z = 402 M+1$;

30

N-(Benzyloxymethyl-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(1-Cyano-3-phenyl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 426 M+1$;

35

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (2-benzyloxy-1-cyano-ethyl)-amide. MS: $m/z = 416 M+1$;

5 4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (1-cyano-3-phenyl-propyl)-amide. MS: $m/z = 400 M+1$;

2-(Morpholine-4-carbonyl)-cyclohexanecarboxylic acid (benzyloxymethyl-cyano-methyl)-amide;

10 *N*-(2-Benzyloxy-1-cyano-ethyl)-4-morpholin-4-yl-2-naphthalen-2-ylmethyl-4-oxo-butyramide. MS: $m/z = 486 M+1$;

N-[2-(4-Chloro-benzyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 477 M+1$;

15 *N'*-(Benzyloxymethyl-cyano-methyl)-*N'*-[4-(5-chloro-H-benzoimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamide;

20 *N'*-[2-(4-Acetylamino-phenyl)-*N'*-(2-benzyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-succinamide. MS: $m/z = 505 M+1$;

N-(cyano-1-methyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*-(cyano-cyclopropyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide. MS: $m/z = 407 M+1$;

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (1-benzyl-4-cyano-piperidin-4-yl)-amide. MS: $m/z = 455 M+1$;

5 *N*-(1-Benzyl-4-cyano-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481 M+1$;

N-[1-(3-Benzoyloxy-benzyl)-3-cyano-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 573 M+1$;

10

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$;

15

N-[3-Cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 503 M+1$;

N-[3-Cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 503 M+1$;

20

N-[3-Cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 535 M+1$;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 459 M+1$;

25

4,4-Dimethyl-2-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl]-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 460 M+1$;

30

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 447 M+1$;

4,4-Dimethyl-2-(2-oxo-2-piperidin-1-yl-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 445 M+1$;

5 4,4-Dimethyl-2-(2-oxo-2-thiomorpholin-4-yl-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 463 M+1$;

N-(3-cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-(3-Cyano-1-isopropyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 419 M+1$;

N-(3-Cyano-1-methyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 391 M+1$;

15 *N*-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 447 M+1$;

20 *N*-(3-Cyano-1-isobutyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 433 M+1$;

N-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 431 M+1$;

25 *N*-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481 M+1$;

N-(3-Cyano-1-methyl-piperidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 405 M+1$;

30

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$;

5 *N*-(3-Cyano-1-propyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 419 M+1$;

N-(3-Cyano-1-cyclopentyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 445 M+1$;

10 *N*-(1-cyano-3-piperidin-1-yl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

(4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(1-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

15 *N*-[1-Cyano-3-(cyclohexyl-ethyl-amino)-propyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 475 M+1$;

20 *N*-[3-(Benzyl-isopropyl-amino)-1-cyano-propyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 497 M+1$;

N-[3-Cyano-1-(1*H*-indol-2-ylmethyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*⁴-Carbamoylmethyl-*N*¹-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-*N*⁴-methyl-succinamide;

N-(3-Cyano-1-cycloheptyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$;

N-[3-Cyano-1-(1,2,3,4-tetrahydro-naphthalen-1-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 507 M+1$;

5 *N*-(1-Bicyclo[2.2.1]hept-2-yl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 471 M+1$;

N-(4-Cyano-1-propyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 433 M+1$;

10 *N*-(1-Benzyl-4-cyano-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 487 M+1$;

N-[3-Cyano-1-(tetrahydro-pyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 461 = M+1$;

15 *N*-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 477 = M+1$;

20 *N*-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 477 = M+1$;

N-[3-Cyano-1-(2-phenyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, $m/z 543=M+1$;

25 *N*-[3-Cyano-1-(3-phenyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, $m/z 543=M+1$;

N-{1-[(benzyl-methyl-amino)-methyl]-1-cyano-3-phenyl-propyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30

N-(4-Cyano-1,2-dimethyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

5 *(4R)*-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butylamide;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(*trans*-4-phenyl-cyclohexylmethyl)-butylamide;

10 2-Bicyclo[4.1.0]hept-7-ylmethyl-*N*-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butylamide;

2-Bicyclo[4.1.0]hept-7-ylmethyl-*N*-(4-cyano-1-methyl-piperidin-4-yl)-4-morpholin-4-yl-4-oxo-butylamide;

15 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydro-naphthalen-2-ylmethyl)-butylamide; MS, m/z 507= $M+1$;

20 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-indan-2-ylmethyl-4-morpholin-4-yl-4-oxo-butylamide; MS, m/z 493= $M+1$;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclopentylmethyl-4-morpholin-4-yl-4-oxo-butylamide; MS, m/z 445= $M+1$;

25 [2-Cyano-2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamino)-ethyl]-carbamic acid tert-butyl ester;

N-{Cyano-[(cyclohexyl-ethyl-amino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-{Cyano-[(dibenzylamino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[[{(Benzyl-ethyl-amino)-methyl]-cyano-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(Cyano-piperidin-1-ylmethyl-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 {1-[3-(1-Benzyl-3-cyano-pyrrolidin-3-yl carbamoyl)-4-cyclohexyl-butyryl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester. MS: $m/z = 566.5 M+1$;

{1-[3-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl carbamoyl)-4-cyclohexyl-butyryl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester. MS: $m/z = 558.6 M+1$;

15 N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-dimethylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 494.5 M+1$;

20 N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 544.4 M+1$;

25 N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 536.4 M+1$;

N-[(3S)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[(3R)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3S)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3R)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[1-cis-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[1-cis-(4-tert-Butyl-cyclohexyl)-(3R)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

5 N-[(3R)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3S)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

10 N-[(3R)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

15 N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3R)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

20 N-[(3S)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3R)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

25 N-[(3S)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

30 N-[(3R)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3S)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3R)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[(3R)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(S)-Cyano-methyl-phenyl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(R)-Cyano-methyl-phenyl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(S)-Cyano-methyl-pyridin-4-yl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(R)-Cyano-methyl-pyridin-4-yl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3--Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[(3R)-3--Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide and

the pharmaceutically acceptable salts, esters or tautomers thereof.

15 Of the aforementioned compounds, preferred are the following:

4-Methyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (2-benzyloxy-1-cyano-ethyl)-amide. MS: $m/z = 402\ M+1$;

20 *N*-(Benzyloxymethyl-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*-(1-Cyano-3-phenyl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 426\ M+1$;

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (2-benzyloxy-1-cyano-ethyl)-amide. MS: $m/z = 416\ M+1$;

30 4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (1-cyano-3-phenyl-propyl)-amide. MS: $m/z = 400\ M+1$;

N-(2-Benzoyloxy-1-cyano-ethyl)-4-morpholin-4-yl-2-naphthalen-2-ylmethyl-4-oxo-butylamide. MS: $m/z = 486 M+1$;

5 *N*-[2-(4-Chloro-benzoyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 477 M+1$;

N-(cyano-1-methyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

10

N-(cyano-cyclopropyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

15

N-(1-Benzyl-4-cyano-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 481 M+1$;

N-[1-(3-Benzoyloxy-benzyl)-3-cyano-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 573 M+1$;

20

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 473 M+1$;

25

N-[3-Cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 503 M+1$;

N-[3-Cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 503 M+1$;

30

N-[3-Cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 535 M+1$;

5 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 459 M+1$;

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 447 M+1$;

10 4,4-Dimethyl-2-(2-oxo-2-thiomorpholin-4-yl-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 463 M+1$;

N-(3-cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 *N*-(3-Cyano-1-isopropyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 419 M+1$;

20 *N*-(3-Cyano-1-methyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 391 M+1$;

N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 447 M+1$;

25 *N*-(3-Cyano-1-isobutyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 433 M+1$;

30 *N*-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 431 M+1$;

N-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481 M+1$;

5 *N*-(3-Cyano-1-methyl-piperidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 405 M+1$;

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$;

10 *N*-(3-Cyano-1-propyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 419 M+1$;

N-(3-Cyano-1-cyclopentyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 445 M+1$;

15 (4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(1-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

20 *N*-[1-Cyano-3-(cyclohexyl-ethyl-amino)-propyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 475 M+1$;

N-[3-(Benzyl-isopropyl-amino)-1-cyano-propyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 497 M+1$;

25 *N*-[3-Cyano-1-(1*H*-indol-2-ylmethyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

*N*⁴-Carbamoylmethyl-*N*¹-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-*N*⁴-methyl-succinamide;

30

N-(3-Cyano-1-cycloheptyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$;

5 *N*-[3-Cyano-1-(1,2,3,4-tetrahydro-naphthalen-1-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 507 M+1$;

N-(1-Bicyclo[2.2.1]hept-2-yl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 471 M+1$;

10 *N*-(4-Cyano-1-propyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 433 M+1$;

N-(1-Benzyl-4-cyano-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 487 M+1$;

15 *N*-[3-Cyano-1-(tetrahydro-pyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 461 = M+1$;

20 *N*-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 477 = M+1$;

N-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 477 = M+1$;

25 *N*-[3-Cyano-1-(2-phenyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, $m/z 543=M+1$;

N-[3-Cyano-1-(3-phenyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, $m/z 543=M+1$;

(4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

5 2-Bicyclo[4.1.0]hept-7-ylmethyl-*N*-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butyramide;

2-Bicyclo[4.1.0]hept-7-ylmethyl-*N*-(4-cyano-1-methyl-piperidin-4-yl)-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydro-naphthalen-2-ylmethyl)-butyramide; MS, m/z 507= $M+1$;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-indan-2-ylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 493= $M+1$;

15 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclopentylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 445= $M+1$;

20 [2-Cyano-2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-ethyl]-carbamic acid tert-butyl ester;

N-{Cyano-[(cyclohexyl-ethyl-amino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*-{Cyano-[(dibenzylamino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-{[(Benzyl-ethyl-amino)-methyl]-cyano-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30

N-(Cyano-piperidin-1-ylmethyl-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

{1-[3-(1-Benzyl-3-cyano-pyrrolidin-3-yl carbamoyl)-4-cyclohexyl-butyryl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester. MS: $m/z = 566.5 M+1$;

{1-[3-(3-Cyano-1-cyclohexyl -pyrrolidin-3-yl carbamoyl)-4-cyclohexyl-butyryl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester. MS: $m/z = 558.6 M+1$;

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-dimethylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 494.5 M+1$;

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 544.4 M+1$;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 536.4 M+1$;

N-[(3S)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3R)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[1-cis-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[1-cis-(4-tert-Butyl-cyclohexyl)-(3R)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[(3R)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3R)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30

N-[(3S)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3R)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[(3R)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3R)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3--Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3R)-3--Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-Cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide and

N-[(3R)-3-Cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.

25 Of the above-listed preferred compounds, the following are more preferred compounds of the invention:

30 *N*-(Benzyloxymethyl-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[2-(4-Chloro-benzyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 477 M+1$;

5 *N*-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 473 M+1$;

10 *N*-[3-Cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 503 M+1$;

N-[3-Cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 503 M+1$;

15 *N*-[3-Cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 535 M+1$;

20 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 459 M+1$;

N-(3-cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

25 *N*-(3-Cyano-1-isopropyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 419 M+1$;

N-(3-Cyano-1-isobutyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. MS: $m/z = 433 M+1$;

30

N-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 431\text{ M}+1$;

5 *N*-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481\text{ M}+1$;

N-(3-Cyano-1-methyl-piperidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 405\text{ M}+1$;

10 *N*-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473\text{ M}+1$;

N-(3-Cyano-1-propyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 419\text{ M}+1$;

15 *N*-(3-Cyano-1-cyclopentyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 445\text{ M}+1$;

20 *N*-[3-Cyano-1-(1*H*-indol-2-ylmethyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-cycloheptyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473\text{ M}+1$;

25 *N*-[3-Cyano-1-(1,2,3,4-tetrahydro-naphthalen-1-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 507\text{ M}+1$;

N-(1-Bicyclo[2.2.1]hept-2-yl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 471\text{ M}+1$;

30

N-[3-Cyano-1-(tetrahydro-pyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: m/z 461 = M+1;

5 N-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: m/z 477 = M+1;

N-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: m/z 477 = M+1;

10 (4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

2-Bicyclo[4.1.0]hept-7-ylmethyl-*N*-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butyramide;

15 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydro-naphthalen-2-ylmethyl)-butyramide; MS, m/z 507=M+1;

20 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-indan-2-ylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 493=M+1;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclopentylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 445=M+1;

25 *N*-{Cyano-[(cyclohexyl-ethyl-amino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(Cyano-piperidin-1-ylmethyl-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30

N-[(3S)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3S)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3S)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3S)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30

N-[(3S)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

5 N-[(3S)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3R)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

10 N-[(3S)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3S)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

15 N-[(3S)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

20 N-[(3S)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3S)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

25 N-[(3S)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

30 N-[(3S)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide;

N-[(3R)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3S)-3-Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide and

N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.

10 Of the above-listed more preferred compounds, the following are most preferred compounds of the invention:

N-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 459 M+1$;

N-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 431 M+1$;

20

N-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481 M+1$;

25

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$;

(4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

30

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydro-naphthalen-2-ylmethyl)-butyramide; MS, m/z 507=M+1;

5 *N*-[(3S)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-[(3R)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide and

N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.

15

Any compounds of this invention containing one or more asymmetric carbon atoms may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. All such isomeric forms of these compounds are expressly
20 included in the present invention. Each stereogenic carbon may be in the R or S configuration, or a combination of configurations.

Some of the compounds of formulas (I) and (Ia) can exist in more than one tautomeric form. The invention includes all such tautomers.

25

It shall be understood by one of ordinary skill in the art that all compounds of the invention are those which are chemically stable.

The invention includes pharmaceutically acceptable derivatives of compounds of
30 formulas (I) and (Ia). A "pharmaceutically acceptable derivative" refers to any pharmaceutically acceptable salt or ester of a compound of this invention, or any other

compound which, upon administration to a patient, is capable of providing (directly or indirectly) a compound of this invention, a pharmacologically active metabolite or pharmacologically active residue thereof.

- 5 In addition, the compounds of this invention include prodrugs of compounds of the formulas (I) and (Ia). Prodrugs include those compounds that, upon simple chemical transformation, are modified to produce the compounds of the invention. Simple chemical transformations include hydrolysis, oxidation and reduction. Specifically, when a prodrug of this invention is administered to a patient, the prodrug may be transformed
10 into a compound of formula (I) and (Ia), thereby imparting the desired pharmacological effect.

In order that the invention herein described may be more fully understood, the following detailed description is set forth. As used herein, the following abbreviations are used:

15

BOC or t-BOC is tertiary butoxycarbonyl

t-Bu is tertiary butyl

DMF is dimethylformamide

EtOAc is ethyl acetate

20

THF is tetrahydrofuran

Ar is argon

EDC is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

HOBT is 1-hydroxybenzotriazole.

25

Also, as used herein, each of the following terms, used alone or in conjunction with other terms, are defined as follows (except where noted to the contrary):

30

The term "alkyl" refers to a saturated aliphatic radical containing from one to ten carbon atoms or a mono- or polyunsaturated aliphatic hydrocarbon radical containing from two to twelve carbon atoms, containing at least one double or triple bond, respectively. "Alkyl" refers to both branched and unbranched alkyl groups. Preferred alkyl groups are

straight chain alkyl groups containing from one to eight carbon atoms and branched alkyl groups containing from three to eight carbon atoms. More preferred alkyl groups are straight chain alkyl groups containing from one to six carbon atoms and branched alkyl groups containing from three to six carbon atoms. It should be understood that any combination term using an "alk" or "alkyl" prefix refers to analogs according to the above definition of "alkyl". For example, terms such as "alkoxy", "alkythio" refer to alkyl groups linked to a second group via an oxygen or sulfur atom. "Alkanoyl refers to an alkyl group linked to a carbonyl group (C=O).

10 The term "cycloalkyl" refers to the cyclic analog of an alkyl group, as defined above, optionally unsaturated and substituted. Preferred cycloalkyl groups are saturated cycloalkyl groups containing from three to eight carbon atoms, and more preferably three to six carbon atoms.

15 The term "aryl" refers to phenyl and naphthyl.

Each of the above defined "alkyl", "cycloalkyl" and "aryl" shall be understood to include their halogenated analogs.

20 The term "halo" refers to a halogen radical selected from fluoro, chloro, bromo or iodo. Preferred halo groups are fluoro, chloro and bromo.

The term "heteroaryl" refers to a stable 5-8 membered (but preferably, 5 or 6 membered) monocyclic or 8-11 membered bicyclic aromatic heterocycle radical. Each heterocycle consists of carbon atoms and from 1 to 4 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. The heterocycle may be attached by any atom of the cycle, which results in the creation of a stable structure. Preferred heteroaryl radicals include, for example, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzothienyl, indazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, purinyl,

quinoliziny, quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl or phenoxazinyl,

- 5 The term "heterocycle" refers to a stable 5-8 membered (but preferably, 5 or 6 membered) monocyclic or 8-11 membered bicyclic heterocycle radical which may be either saturated or unsaturated, and is non-aromatic. Each heterocycle consists of carbon atoms and from 1 to 4 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. The heterocycle may be attached by any atom of the cycle, which
- 10 results in the creation of a stable structure. Preferred heterocycle radicals include, for example, pyrrolinyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, pyranyl, thiopyranyl, piperazinyl, indolinyl, azetidyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuran, hexahydropyrimidinyl, hexahydropyridazinyl, 1,4,5,6-tetrahydropyrimidin-2-ylamine, dihydro-oxazolyl, 1,2-
- 15 thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide and imidazolidinyl-2,4-dione.

- The terms "heterocycle", "heteroaryl" or "aryl", when associated with another moiety, unless otherwise specified shall have the same meaning as given above. For example,
- 20 "aroyl" refers to phenyl or naphthyl linked to a carbonyl group (C=O).

- Each aryl or heteroaryl unless otherwise specified includes its partially or fully hydrogenated derivative. For example, quinolinyl may include decahydroquinolinyl and tetrahydroquinolinyl, naphthyl may include its hydrogenated derivatives such as
- 25 tetrahydronaphthyl. Other partially or fully hydrogenated derivatives of the aryl and heteroaryl compounds described herein will be apparent to one of ordinary skill in the art.

- As used herein above and throughout this application, "nitrogen" and "sulfur" include any oxidized form of nitrogen and sulfur and the quaternized form of any basic nitrogen.

30

In order that this invention be more fully understood, the following examples are set forth. These examples are for the purpose of illustrating preferred embodiments of this invention, and are not to be construed as limiting the scope of the invention in any way.

- 5 The examples which follow are illustrative and, as recognized by one skilled in the art, particular reagents or conditions could be modified as needed for individual compounds. Starting materials used in the scheme below are either commercially available or easily prepared from commercially available materials by those skilled in the art.

10

GENERAL SYNTHETIC METHODS

The invention also provides processes of making the present novel compounds.


- 15 Compounds of the invention may be prepared by methods described below. Standard peptide coupling, protection and deprotection reactions (see for example M. Bodanszky, The Practice of Peptide Synthesis, Springer-Verlag, 1984) are employed in these syntheses. Alkylated succinic acid derivatives used as starting materials are either commercially available or easily prepared by methods known to those skilled in the art.
- 20 For example succinate derivatives may be alkylated by treatment with a suitable alkylating agent such as an alkyl halide in the presence of a suitable base such as lithium diisopropyl amide. If the succinate derivative contains a chiral auxiliary, for example a succinic acid chiral oxazolidine amide, a chiral alkylated succinate derivative can be obtained as described by Azam et al. (J. Chem. Soc. Perkin Trans. 1, 1996, 621) and
- 25 Evans et al. (J. Am. Chem. Soc., 1981, 103, 2127). Alternatively, racemic alkylated succinate derivatives may be resolved by methods known to those skilled in the art if desired, for example by selective enzymatic hydrolysis as described by Oikawa et al. (Tetrahedron Lett., 1996, 37, 6169), Wirz et al. (Tetrahedron Asymmetry, 1997, 8, 187) and Ozaki et al. (Chem. Lett., 1997, 741).

30

Compounds of the invention in which R_1 is an amino group or a heterocyclyl group containing a nitrogen that forms an amide bond with A ($A = -C(O)-$) of formula I may be prepared as described below and illustrated in Scheme I.

5

Scheme I



10


A succinic acid monoester, for example a methyl ester as shown, is reacted with the desired amine R_1H under standard peptide coupling conditions. An example of standard coupling conditions would be combining the starting materials in the presence of a coupling reagent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) with 1-hydroxybenzotriazole (HOBT), in a suitable solvent such as DMF or methylene chloride. A base such as N-methylmorpholine may be added. The resulting amide ester is hydrolyzed in aqueous base or acid. A suitable co-solvent such as THF or MeOH may be added. The resulting acid is then coupled with an amino nitrile under standard peptide coupling conditions as described above to provide the desired compound of formula (I).

15

Alternately, one may couple the acid with an amino amide and then react the resulting amide with a suitable dehydrating agent such as cyanuric chloride in a suitable solvent such as DMF to provide the desired nitrile of formula (I).

- 5 Compounds of the invention in which R_1 is alkyl or cycloalkyl may be prepared by reacting a succinic acid ester derivative, for example a methyl ester as shown below, containing an activated amide, for example a N-methoxy-N-methylamide (Solladic-Cavallo et al., *Tetrahedron Asymmetry*, 1996, 7, 1797), or a N-imidazolium-N-methylamide (De Las Heras et al., *Tetrahedron Lett.*, 1997, 38, 1817) with a suitable
10 organometallic reagent, such as an alkylmagnesium bromide (Scheme II). The resulting ketoester is then carried forth as described for the amide ester intermediate in Scheme I.

Scheme II




15

- Compounds of the invention in which R_1 is aryl or heteroaryl may be prepared by reacting a suitable aryl or heteroaryl organometallic reagent with an activated amide as described above for R_1 = alkyl or cycloalkyl. In addition, an acyl halide, for example an acid chloride, may be used in place of an activated amide. Suitable aryl or heteroaryl
20 organometallic reagents for reaction with acyl halides are known in the art and include aryl magnesium bromides (F. Babudri et al., *Tetrahedron*, 1996, 52, 13513), aryl lithium reagents in the presence of ZnCl_2 (C. Kim et al., *Tetrahedron Lett.*, 1994, 35, 3017) and aryl stannanes under Pd catalyzed Stille coupling conditions (M.J. Plunkett et al., *J. Am. Chem. Soc.*, 1995, 117, 3306).

25

A method which may be used to stereoselectively prepare alkylated succinic acid derivatives, useful in the procedures described in Schemes I and II, is illustrated in Scheme III.

5 Scheme III



In this procedure, a carboxylic acid bearing R_4 and R_5 is coupled with a chiral auxiliary group known in the art (see for example D. Evans *et al.*, J. Am. Chem. Soc., 1990, 4011; D. Evans *et al.*, J. Am. Chem. Soc., 1981, 2127, 103; D. Evans *et al.*, Tetrahedron Lett, 1987, 6141; D. Evans *et al.*, Tetrahedron Lett, 1987, 1123), such as (*S*)-(-)-4-isopropyl-2-oxazolidinone as illustrated in Scheme III. Suitable coupling conditions would include first converting the carboxylic acid to an acid chloride, for example by reaction with oxalyl chloride and DMF in a suitable solvent such as methylene chloride at about 0 °C to room temperature. The acid chloride may then be coupled to the oxazolidinone in a suitable solvent, such as THF, in the presence of a suitable base, such as *n*-BuLi at about -78 °C to room temperature.

The resulting product is then alkylated with a halomethylcarbonyl compound bearing R₁, R₂ and R₃ in a suitable solvent such as THF, in the presence of a suitable base, such as sodium bis(trimethylsilyl)amide at about -78 °C to room temperature. The large group on the chiral auxiliary (the isopropyl group in the illustrated example) directs the alkylating group to the side opposite it providing predominantly one isomer at the alkylated carbon. The chiral auxiliary is then removed by methods known in the art, for example in the case illustrated, in a suitable solvent such as THF in the presence of water and a suitable peroxide such as 30% hydrogen peroxide at about 0 °C, providing the desired succinic acid derivative which may be used in the procedure outlined in Scheme I, or converted to the corresponding ester by methods known to those skilled in the art and used in the procedure outlined in Scheme II.


Compounds of the invention in which R₁ is alkyl, cycloalkyl, aryl or heteroaryl and A is -C(O)- may be reacted with a suitable reducing agent such as sodium borohydride to produce the corresponding compound where A is -C(OH)-

The synthetic examples below are illustrative of the methods used to prepare the compounds of the invention.

SYNTHETIC EXAMPLES

EXAMPLE 1

N-(Benzyloxymethyl-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide



(a) (R)-2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butanoic acid 1-methyl ester

(R)-2-(Cyclohexylmethyl) succinic acid 1-methyl ester (1.00 g, 4.38 mmol) was dissolved in 10 mL of DMF and cooled to 0 °C with an ice-water bath. EDC (1.12 g, 5.69 mmol) then 1-hydroxybenzotriazole (0.77 g, 5.69 mmol) were added and stirring continued, under argon, for 25 min. Morpholine (0.76 mL, 8.76 mmol) was added and stirring was continued overnight (16 h). The solution was diluted with 200 mL of EtOAc and washed with a 1.0 M solution of HCl (3 x 100 mL), a saturated solution of NaHCO₃ (3 x 100 mL), water (100 mL), then with brine (100 mL). The organic layer was dried over MgSO₄, filtered and concentrated by rotary evaporation to give (R)-2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butanoic acid 1-methyl ester (1.26 g, 99%) as a white solid. The product was used in the following step without purification.

(b) (R)-2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butanoic acid

The ester from the above step (1.21 g, 4.06 mmol) was dissolved in THF (55 mL) and MeOH (27 mL) was then added. This solution was cooled to 0 °C with an ice-water bath, then a solution of LiOH·H₂O (0.51 g, 12.18 mmol) in H₂O (27 mL) was slowly added and the mixture was allowed to stir for 5 hr at room temperature. The mixture

was poured into water (100 mL) and extracted with Et₂O (2 x 50 mL). The aqueous layer was acidified with 1.0 N HCl, then extracted with EtOAc (3 x 50 mL). The combined organic layers were then washed with H₂O (2 x 50 mL) followed by brine (2 x 50 mL), dried over Na₂SO₄, filtered and the solvent was evaporated to afford (R)-2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyric acid (0.88 g, 76%) as a clean white solid that was used without further purification.

(c) *N*-(*t*-butoxycarbonyl)-L-(O-benzyl)serinamide

NH₄OH (10 mL) was added to a premixed (15 min) solution of *N*-(*t*-butoxycarbonyl)-L-(O-benzyl)serine (10.0 g, 33.9 mmol), EDC (7.80 g, 40.7 mmol), and HOBT (5.50 g, 40.7 mmol) in DMF (120 mL) at room temperature. After 16 h the reaction mixture was diluted with CH₂Cl₂ and filtered, washed sequentially with 10% aq. HCl, satd. NaHCO₃, H₂O (x3), brine, then dried over Na₂SO₄, and concentrated giving *N*-(*t*-butoxycarbonyl)-L-(O-benzyl)serinamide (11.5 g) as a white solid.

(d) L-(O-Benzyl) Serine amide HCl

N-(*t*-butoxycarbonyl)-L-(O-benzyl)serinamide (1.00 g) was dissolved in anhydrous THF (5 ml) and cooled in an ice/water bath. A 4M solution of HCl in dioxane (5 ml) was added and the reaction stirred for two h and then concentrated under reduced pressure to provide a glass which was then triturated with diethyl ether to obtain L-(O-Benzyl) Serine amide HCl as a white solid (0.9 g). This material was used without further purification.

(e) *N*-(2-Benzyl-1-carbamoyl-ethyl)-2-cyclohexylmethyl-4-morpholin-4-yl-oxo-butyramide



(R)-2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyric acid (0.20 g, 0.71 mmol) was dissolved in 6 mL of DMF and cooled to 0 °C with an ice-water bath. EDC (0.18 g, 0.92 mmol) and 1-hydroxybenzotriazole (0.12 g, 0.92 mmol) were added and stirring, under argon, continued for 25 min. To the cold solution was added the HCl salt of O-benzyl-L-serineamide (0.16 g, 0.71 mmol), followed by addition of *N*-methylmorpholine (0.23 mL, 2.10 mmol) and stirring was continued overnight (16 h). The solution was diluted with 100 mL of EtOAc and washed with a 1.0 M solution of HCl (3 x 10 mL), a saturated solution of NaHCO₃ (3 x 10 mL), water (100 mL) then

with brine (100 mL). The organic layer was dried over MgSO_4 , filtered and concentrated by rotary evaporation to give (0.34 g, 106%) of the crude product which was chromatographed (SiO_2 , 2% MeOH in EtOAc) to give N-(2-Benzyloxy-1-carbamoyl-ethyl)-2-cyclohexylmethyl-4-morpholin-4-yl-oxo-butyramide (0.22 g, 69%).


(f) N-(Benzyloxymethyl-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-(2-Benzyloxy-1-carbamoyl-ethyl)-2-cyclohexylmethyl-4-morpholin-4-yl-oxo-

butyramide (0.22 g, 0.48 mmol) was dissolved in 4 mL of DMF and cooled to 0 °C with an ice-water bath. To this solution was added cyanuric chloride (0.089 g, 0.48 mmol) and the reaction mixture was stirred for one h. The reaction was quenched with cold water (2 mL), extracted with EtOAc (3 x 100 mL), and the organic layers were combined and washed with 100 mL brine. The organic layer was dried over MgSO_4 , filtered and concentrated by rotary evaporation to give the crude residue. The residue was purified by chromatography (SiO_2 , 35% hexane in EtOAc) to give the title compound as a clear oil (0.12 g, 57%); ^1H NMR (CDCl_3) δ 7.35 (5H, m), 7.02-7.08 (1H, m), 5.01-5.09 (1H, m), 4.52-4.63 (2H, m), 3.72-3.34 (10H, m), 2.96-2.83 (1H, m), 2.78-2.62 (1H, m), 2.18-2.30 (1H, m), 1.75-1.51 (6H, m), 1.30-1.07 (5H, m), 0.95-0.73 (2H, m). MS: m/z = 442 $M+1$.


EXAMPLE 2**2-(Morpholine-4-carbonyl)-cyclohexanecarboxylic acid (benzyloxymethyl-cyano-methyl)-amide****(a) 2-(Morpholine-4-carbonyl)-cyclohexanecarboxylic acid**

To a solution of *cis*-1,2-cyclohexane dicarboxylic anhydride (1.0 g, 6.49 mmol) in CH₂Cl₂ (16 mL) was added morpholine (0.56 g, 6.49 mmol) at 0 °C, the resulting mixture was allowed to stir at room temperature overnight (20 h). When complete the reaction mixture was washed with HCl (1.0 N, 20 mL) then with H₂O (2 x 20 mL). The combined aqueous layers were extracted with CH₂Cl₂ (3 x 10 mL), and the combined organic layers were dried over MgSO₄, the solvent was evaporated in vacuo to give the title compound (1.4 g, 90 %) as a white solid which was used later without further purification.

(b) 2-(Morpholine-4-carbonyl)-cyclohexanecarboxylic acid (2-benzyloxy-1-carbamoyl-ethyl)-amide

This compound was prepared by the procedure described in Example 1 part e) from 2-(Morpholine-4-carbonyl)-cyclohexanecarboxylic acid and the HCl salt of O-benzyl-L-serineamide to give the crude product which was chromatographed (SiO₂, 10% Hexanes in EtOAc) to yield the title compound (1.2 g, 50% yield).


(c) 2-(Morpholine-4-carbonyl)-cyclohexanecarboxylic acid (benzyloxymethyl-cyano-ethyl)-amide




The amide from the step above (0.10 g, 0.24 mmol) and cyanuric chloride (0.044 g, 0.24 mmol) were reacted as described by the procedure in Example 1 part (f). The residue was purified by chromatography (SiO₂, 50% Hexanes in EtOAc) to give the title compound as a clear oil (0.055 g, 60%); ¹H NMR (CDCl₃) 7.38-7.1 (6H, m), 5.0 (1H, m), 4.6 (2H, m), 3.75-3.36 (9H, m), 3.3-3.05 (1H, m), 2.55-2.3 (2H, m), 2.0-1.15 (8H, m). MS: m/z = 400 M+1.


EXAMPLE 3

N¹-(Benzyloxymethyl-cyano-methyl)-N⁴-[4-(5-chloro-H-benzoimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamide




(a) (5-Chloro-2-(4-nitro-phenyl)-1H-benzoimidazole

To a solution of 4-Nitrobenzaldehyde (6.4 g, 42.35 mmol) in THF (300 mL) was added 4-Chloro-1,2-phenylenediamine (4.0 g, 28.05 mmol), and the mixture was stirred under air at room temperature for two days. When complete the reaction mixture was diluted with EtOAc (200 mL), washed with H₂O (200 mL) then with HCl (1.0 N, 200 mL). The product precipitated out and was collected by filtration under suction. The solid was washed with EtOAc and allowed to dry. The product was collected as a brownish solid (4.2 g, 55%), pure HCl salt.

(b) 4-(5-Chloro-1H-benzoimidazol-2-yl)-phenylamine


In 500 mL round bottom flask was suspended 5-Chloro-2-(4-nitro-phenyl)-1H-benzoimidazole (4.23 g, 13.66 mmol) in AcOH (30 mL) at 0 °C, and to the suspension was added dropwise SnCl₂·H₂O (9.4 g, 41.66 mmol), in HCl (15 mL). After the addition, the ice bath was removed and the mixture was allowed to stir at room temperature overnight (16 h). The mixture was poured onto ice, and NaOH (50% aq.) was slowly added to pH 12. Then the mixture was extracted with EtOAc (4 x 150 mL). The combined organic layers were washed with brine, dried over NaSO₄ and filtered, the solvent was evaporated to yield the product as an orange oil. The oil was dissolved in EtOAc and washed with HCl (1.0 N, 100 mL) and the product was isolated as HCl salt (2.13 g, 67%).

(c) *N*-[4-(5-Chloro-1*H*-benzoimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamic acid methyl ester




This compound was prepared by the procedure described in Example 1 part (a) from 4-(5-Chloro-1*H*-benzoimidazol-2-yl)-phenylamine and (R)-2-(Cyclohexylmethyl) succinic acid 1-methyl ester to give the crude product (2.0 g, 51%), which was used in the next step without further purification.

(d) *N*-[4-(5-Chloro-1*H*-benzoimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamic acid




This compound was prepared by the procedure described in Example 1 part (b) from *N*-[4-(5-Chloro-1*H*-benzoimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamic acid methyl ester, and was isolated as an HCl salt.

(e) *N*¹-(2-Benzoyloxy-1-carbamoyl-ethyl)-*N*⁴-[4-(5-chloro-1*H*-benzoimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamide



This compound was prepared by the procedure described in Example 1 part (e) from *N*-[4-(5-Chloro-1*H*-benzimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamic acid and the HCl salt of O-benzyl-L-serineamide to give the crude product which was chromatographed (SiO₂, 30% hexanes in EtOAc) to yield the title compound in 76% yield.


(f) *N*¹-(Benzyloxymethyl-cyano-methyl)-*N*⁴-[4-(5-chloro-*H*-benzimidazol-2-yl)-phenyl]-2-cyclohexylmethyl-succinamide




The amide from the step above (0.08 g, 0.13 mmol) and cyanuric chloride (0.024 g, 0.13 mmol) were reacted as described by the procedure in Example 1 part (f). The residue was purified by chromatography (SiO₂, 35% MeOH in CH₂Cl₂) to give the title compound as two separated diastereomers (0.05 g, 64%); ¹H NMR (CDCl₃) 8.2-7.92 (2H, m), 7.8-7.72 (2H, m), 7.57-7.44 (2H, m), 7.3-7.15 (6H, m), 5.47 (1H, m), 4.5-4.4 (2H, m), 3.65-3.5 (2H, m), 3.1-3.0 (1H, m), 2.67-2.54 (1H, m), 2.47-2.34 (1H, m), 1.95-1.5 (7H, m), 1.4-1.15 (6H, m), 1.0-0.8 (3H, m). MS: *m/z* = 599 *M*+1.


EXAMPLE 4

N-(Cyano-cyclopropyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide




(a) 1-Amino-1-cyclopropanecarboxamide HCl

This compound was prepared by the procedures described in Example 1 part (c and d) from 1-tert-Butoxycarbonylamino-cyclopropanecarboxylic acid.

(b) 1-(2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-cyclopropanecarboxylic acid amide

1-(2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-cyclopropanecarboxylic acid amide was prepared by the procedure described in Example 1 part (e) from (R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyric acid (0.20 g, 0.71 mmol), EDC (0.18 g, 0.92 mmol), 1-hydroxybenzotriazole (0.12 g, 0.92 mmol) and the HCl salt of 1-amino-1-cyclopropanecarboxamide (0.096 g, 0.71 mmol) to yield a white solid (0.084, 33%) after purification by preparative HPLC.


(c) N-(Cyano-cyclopropyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

The amide from the step above (0.084 g, 0.23 mmol) and cyanuric chloride (0.042 g, 0.23 mmol) were reacted as described by the procedure in Example 1 part (f). The residue was purified by chromatography (SiO₂, 2% MeOH in CH₂Cl₂) to give the title compound as a clear oil (0.03 g, 38%); ¹H NMR (CDCl₃) 7.38 (1H, m), 3.75-3.36 (8H, m), 2.85-2.68 (2H,


m), 2.26-2.17 (1H, m), 1.74-1.53 (6H, m), 1.52-1.40 (2H), 1.27-1.02 (7H, m), 0.90-0.74 (2H, m). MS: $m/z = 348 M+1$.

EXAMPLE 5

N-(Cyano-1-methyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide




(a) 4-Amino-4-cyano-1-methylpiperidine





A solution of ammonium chloride (1.89 g, 35.37 mmol) and potassium cyanide (2.30 g, 35.37 mmol) was prepared in 50 mL of water. 1-Methyl-4-piperidone (1 g, 8.84 mmol) was added to the solution and stirring was continued for 2 days. The solution was brought to pH 11 with solid sodium carbonate and the reaction solution was extracted with 3 x 100 mL of EtOAc. The organic layer was dried over Na_2SO_4 , decanted and concentrated to an orange oil (857 mg). ^1H NMR showed that the oil was a 2:1:1 mixture of 4-amino-4-cyano-1-methylpiperidine, the corresponding cyanohydrin and starting ketone. The crude mixture was used in the next step without further purification.

(b) N-(Cyano-1-methyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide




(R)-2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butanoate (0.20 g, 0.71 mmol) (see Example 1-part a and b) was dissolved in 5 mL of DMF and cooled to 0 °C by an ice-water bath. EDC (0.18 g, 0.92 mmol) and 1-hydroxybenzotriazole (0.12 g, 0.92 mmol) were added and stirring, under argon, continued for 25 min. Then 4-amino-4-cyano-1-methylpiperidine (0.098 g, of the mixture of aminonitrile:cyanohydrin:ketone) was dissolved in 1 mL of DMF and added to the solution of the active ester. The resulting mixture was stirred at ambient temperature for 4 h. The volatiles were removed in vacuo and the resulting residue was dissolved in 100 mL of EtOAc and washed sequentially with 2x100 mL of saturated sodium bicarbonate and 1x100 mL of brine. The organic layer was dried over MgSO₄, filtered, and concentrated to a thick oil. The oil was purified by column chromatography on SiO₂ using as eluent 5% MeOH in CH₂Cl₂ to give the title compound as a clear oil (0.005 g): ¹H NMR (CDCl₃) 6.64 (1H, m), 3.73-3.37 (8H, m), 2.95-2.85 (1H, m), 2.75-2.55 (2H, m), 2.5-2.37 (2H, m), 2.34-2.25 (3H, m), 1.98-1.82 (2H, m), 1.80-1.58 (6H, m), 1.32-1.10 (8H, m), 0.95-0.78 (3H, m). MS: m/z = 405 M+1.

EXAMPLE 6

N-(Cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide**(a) 3-Amino-3-cyano-1-benzylpyrrolidine**


3-Amino-3-cyano-1-benzylpyrrolidine was prepared by a method analogous to that of Example 5-part (a) with the exception that no sodium carbonate was added to the reaction mixture. The product was extracted from the crude reaction with 3x100 mL of EtOAc and was used without purification.

(b) N-(Cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide. Separated diastereomers


Diastereomeric *N*-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide was prepared by a method analogous to that of Example 3-part (d). The purification was done by reverse phase preparative HPLC to separate the two diastereomers. ¹H NMR (CDCl₃) 7.31 (5H, m), 6.81 (1H, s), 3.66-3.47 (10H, m), 3.21-3.18 (1H, m), 2.85-2.67 (5H, m), 2.3-2.1 (3H, m), 1.72-1.65 (5H, m), 1.25-1.16 (5H, m), 0.95-0.78 (3H, m). MS, m/z 467 = M+1.

EXAMPLE 7

N-(3-Cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide




(a) 1-(3,3-Dimethyl-butyl)-pyrrolidin-3-ol



Acetic acid (31 μ L), pyrrolidin-3-ol (3.47 g, 39.8 mmol) and 3,3-dimethyl-butyraldehyde (3.99 g, 39.8 mmol) were dissolved in THF (100 mL) and stirred at room temperature for 40 min. The solution was then cooled to 0 °C on an ice bath. To the cold solution sodium


triacetoxyborohydride (10.98 g, 51.8 mmol) was added and the mixture was stirred at room temperature overnight. The reaction mixture was basified with a 5.0 M solution of NaOH and extracted with CH_2Cl_2 (3X150 mL). The organic layer was dried over Na_2SO_4 , filtered and concentrated to dryness to afford 1-(3,3-dimethyl-butyl)-pyrrolidin-3-ol in (6.34 g, 92.5 %), which was used without purification.

(b) 1-(3,3-Dimethyl-butyl)-pyrrolidin-3-one




To a dry 500 mL round bottom flask was added oxalyl chloride (9.40 g, 74.06 mmol) in anhydrous CH_2Cl_2 (180 mL) and this solution was cooled to -78°C . DMSO (11.57 g, 148.08 mmol) was added via syringe over a 15 min period during which vigorous gas evolution was observed and the mixture was stirred for 25 min. To the mixture was added 1-(3,3-dimethyl-butyl)-pyrrolidin-3-ol (6.34 g, 37.01 mmol) in CH_2Cl_2 (20 mL) via syringe over a 10 min period. The mixture was stirred for one h at -78°C , then Et_3N (29.96 g, 296.08 mmol) was added via syringe and the dry ice/acetone bath was removed to allow the reaction mixture to warm up to room temperature over 2 h.

The reaction mixture was poured into H_2O (250 mL), the layers were separated and the aqueous was washed with CH_2Cl_2 (3x150 mL). The organic layers were combined and washed with NaHCO_3 (200 mL), brine (200 mL), dried over Na_2SO_4 , filtered, then concentrated. The product was purified by column chromatography on SiO_2 using as eluent 100% CH_2Cl_2 to afford the title compound in (3.5 g, 56 % yield).

(c) 3-Amino-1-(3,3-dimethyl-butyl)-pyrrolidine-3-carbonitrile

To a dry 250 mL round bottom flask set with a condenser was added sodium cyanide (0.786 g, 16.03 mmol), ammonium chloride (0.904 g, 16.9 mmol) and NH_3/MeOH (2.0 M, 15.36 mL, 30.7 mmol). To the mixture was added 1-(3,3-dimethyl-butyl)-pyrrolidin-3-one (2.60 g, 15.36 mmol) and MgSO_4 (2.5 g). The mixture was heated at 60 °C in an oil bath and stirred for 4 h. The mixture was cooled to room temperature and filtered to remove the insolubles, the filtrate was concentrated to dryness and the residue was dissolved in $\text{EtOAc}/\text{Et}_2\text{O}$ and filtered through a pad of diatomaceous earth then concentrated to dryness to give the title compound in (1.5 g, 51% yield). The crude compound was used in the next step without further purification.


(d) *N*-(3-Cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

Diastereomeric *N*-(3-cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide was prepared by a method analogous to that of Example 3-part (d). The purification was done by reverse phase preparative HPLC to separate the two diastereomers. ^1H NMR (CDCl_3) 6.95 (1H, s), 3.7-3.4 (8H, m), 3.1-3.07 (1H,


m), 2.9-2.52 (5H, m), 2.45-2.39 (2H, m), 2.3-2.08 (3H, m), 1.78-1.6 (5H, m), 1.38-1.3 (2H, m), 1.3-1.1 (5H, m), 0.95-0.78 (12H, m). MS, m/z 461 = $M+1$.

EXAMPLE 8


N-(1-cyano-3-piperidin-1-yl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide




(a) *2-tert*-Butoxycarbonylamino-*N*-methoxy-*N*-methyl succinamic acid benzyl ester



This compound was prepared by a method analogous to that of Example 3-part (d) from *N,O*-Dimethylhydroxylamine hydrochloride and 2-*tert*-Butoxycarbonylamino-succinic acid 1-benzyl ester.


(b) 2-*tert*-Butoxycarbonylamino-4-oxo-butyrlic acid benzyl ester

To 2-*tert*-butoxycarbonylamino-*N*-methoxy-*N*-methyl succinamic acid benzyl ester (0.50 g, 1.36 mmol) in THF (5 mL) at -78°C was added DIBAL-H (0.233 g, 1.67 mmol) dropwise. The mixture was stirred for 1.5 h and monitored by TLC. When completed the reaction was quenched with 5 mL of saturated solution of Rochell salt ($\text{KNaC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$) (potassium sodium tartrate) and extracted with EtOAc (2x10 mL). The organic layer was washed with H_2O , brine, dried over Na_2SO_4 , filtered and concentrated in vacuo to give the title compound in quantitative yield (0.40 g).

(c) 2-*tert*-Butoxycarbonylamino-4-piperidin-1-yl-butyrlic acid benzyl ester


This compound was prepared by a method analogous to that of Example 5-part (a) from 2-*tert*-butoxycarbonylamino-4-oxo-butyrlic acid benzyl ester and piperidine. The product was dissolved in 20 % TFA in CH_2Cl_2 (10 mL) at 0°C , then it was allowed to stir for 1 h at room temperature. The solvent was evaporated to give the diamine TFA salt.

(d) 2-(2-Cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-4-piperidin-1-yl-butyr-
ic acid benzyl ester




This compound was prepared by a method analogous to that of Example 3-part (d) from the above TFA diamine salt and (R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyric acid.

(e) *N*-(1-Carbamoyl-3-piperidin-1-yl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide



To 2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-4-piperidin-1-yl-butyr-ic acid benzyl ester (0.10 g, 0.18 mmol) was added NH_3/MeOH (2.0 M, 10 mL) in a sealed tube and the mixture was stirred at room temperature for 3 days. The solvent was evaporated to give the title compound in 50% yield.


- (f) *N*-(1-Cyano-3-piperidin-1-yl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide




This compound was prepared by a method analogous to that of Example 1-part (f) from *N*-(1-Carbamoyl-3-piperidin-1-yl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide. ¹H NMR (CDCl₃) 4.98-4.94 (1H, m), 3.74-3.45 (8H, m), 3.05-2.2 (9H, m), 2.15-1.4 (11H, m), 1.35- 1.05 (7H, m), 0.92-0.75 (3H, m). MS: m/z = 433 M+1.

EXAMPLE 9

N-[3-Cyano-1-(1*H*-indol-2-ylmethyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide




(a) 1-(Toluene-4-sulfonyl)-1*H*-indole-3-carboxaldehyde



This compound was prepared according to the reference below:


Chauhan, P. M. S., Chatterjee, R. K. Indian J. Chem. Sect. B, 1994, 33 (1), 32-37.

(b) *N*-{3-Cyano-1-[1-(toluene-4-sulfonyl)-1*H*-indole-3-ylmethyl]-pyrrolidin-3-yl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide



This compound was prepared by a method analogous to that of Example 3-part (a through d) from 1-(toluene-4-sulfonyl)-1*H*-indole-3-carboxaldehyde and (R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyric acid.

(c) *N*-{3-Cyano-1-(1*H*-indol-2-ylmethyl)-pyrrolidin-3-yl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide



A suspension of *N*-{3-cyano-1-[1-(toluene-4-sulfonyl)-1*H*-indole-3-ylmethyl]-pyrrolidin-3-yl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide (0.052 g, 0.079 mmol) and Mg (powder) (0.038 g, 1.6 mmol) in MeOH (5 mL) was sonicated at room temperature reaction was monitored by MS. When completed, the reaction mixture was poured into a saturated solution of NH₄Cl and extracted with CH₂Cl₂. The combined organic extracts were dried over Na₂SO₄ and concentrated to provide the title compound in 76% yield. The compound was recrystallized from Et₂O/hexanes. ¹H NMR (CDCl₃) . 8.1 (1H, s), 7.71-7.69 (1H, m), 7.39-7.37 (1H, m), 7.25-7.15

(3H, m), 6.9 (1H, s), 4.0-3.8 (2H, m), 3.7-3.3 (1H, m), 3.2-3.1 (1H, m), 3.0-2.5 (3H, m), 2.3-2.15 (2H, m), 1.8-1.4 (5H, m), 1.35-1.03 (5H, m), 0.95-0.78 (3H, m). MS, m/z 506 = $M+1$.

Listed below are additional examples of compounds of the invention, which were prepared by methods analogous to these described above.

4-Methyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (2-benzyloxy-1-cyano-ethyl)-amide.
MS: m/z = 402 $M+1$

N-(1-Cyano-3-phenyl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS:
 m/z = 426 $M+1$

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (2-benzyloxy-1-cyano-ethyl)-amide. MS: m/z = 416 $M+1$.

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (1-cyano-3-phenyl-propyl)-amide. MS: m/z = 400 $M+1$.

N-(2-Benzyloxy-1-cyano-ethyl)-4-morpholin-4-yl-2-naphthalen-2-ylmethyl-4-oxo-butyramide.
MS: m/z = 486 $M+1$.

N-[2-(4-Chloro-benzyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: m/z = 477 $M+1$.

*N*⁴-[2-(4-Acetylamino-phenyl)-*N*¹-(2-benzyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-succinamide. MS: m/z = 505 $M+1$.

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (4-cyano-1-propyl-piperidin-4-yl)-amide. MS: m/z = 407 $M+1$.

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (1-benzyl-4-cyano-piperidin-4-yl)-amide. MS: $m/z = 455 M+1$.

N-(1-Benzyl-4-cyano-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481 M+1$.

N-[1-(3-Benzoyloxy-benzyl)-3-cyano-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 573 M+1$.

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$.

N-[3-Cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 503 M+1$.

N-[3-Cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 503 M+1$.

N-[3-Cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 535 M+1$.

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 459 M+1$.

4,4-Dimethyl-2-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethyl]-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 460 M+1$.

4,4-Dimethyl-2-(2-morpholin-4-yl-2-oxo-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 447 M+1$.

4,4-Dimethyl-2-(2-oxo-2-piperidin-1-yl-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 445 M+1$.

4,4-Dimethyl-2-(2-oxo-2-thiomorpholin-4-yl-ethyl)-pentanoic acid (3-cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-amide. MS: $m/z = 463 M+1$.

N-(3-Cyano-1-isopropyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 419 M+1$.

N-(3-Cyano-1-methyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 391 M+1$.

N-[3-Cyano-1-(1-ethyl-propyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 447 M+1$.

N-(3-Cyano-1-isobutyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 433 M+1$.

N-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 431 M+1$.

N-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 481 M+1$.

N-(3-Cyano-1-methyl-piperidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 405 M+1$.

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$.

N-(3-Cyano-1-propyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.
MS: $m/z = 419 M+1$.

N-(3-Cyano-1-cyclopentyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 445 M+1$.

N-[1-Cyano-3-(cyclohexyl-ethyl-amino)-propyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 475 M+1$.

N-[3-(Benzyl-isopropyl-amino)-1-cyano-propyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 497 M+1$.

N-(3-Cyano-1-cycloheptyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 473 M+1$.

N-[3-Cyano-1-(1,2,3,4-tetrahydro-naphthalen-1-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 507 M+1$.

N-(1-Bicyclo[2.2.1]hept-2-yl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide. MS: $m/z = 471 M+1$.

N-(4-Cyano-1-propyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.
MS: $m/z = 433 M+1$.

N-(1-Benzyl-4-cyano-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.
MS: $m/z = 487 M+1$.

N-[3-Cyano-1-(tetrahydro-pyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: $m/z 461 = M+1$. (1:1 mixture of diastereomers)

N-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: m/z 477 = M+1. (single diastereomer)

N-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: m/z 477 = M+1 (single diastereomer)

N-[3-Cyano-1-(2-phenyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 543=M+1

N-[3-Cyano-1-(3-phenyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 543=M+1

N-4-Cyano-2-phenethyl-2H-pyrazol-3-yl)-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-(4-Cyano-1-phenethyl-1H-pyrazol-3-yl)-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(S)-Cyano-methyl-phenyl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(R)-Cyano-methyl-phenyl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(S)-Cyano-methyl-pyridin-4-yl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(R)-Cyano-methyl-pyridin-4-yl-methyl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3--Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3--Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3-Cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

*N*⁴-Carbamoylmethyl-*N*¹-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-*N*⁴-methyl-succinamide: C₂₅H₄₁N₅O₃, ESMS: 460 (M+1)

{1-[3-(1-Benzyl-3-cyano-pyrrolidin-3-yl carbamoyl)-4-cyclohexyl-butyryl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester. MS: m/z = 566.5 M+1

{1-[3-(3-Cyano-1-cyclohexyl -pyrrolidin-3-yl carbamoyl)-4-cyclohexyl-butyryl]-pyrrolidin-3-yl}-carbamic acid *tert*-butyl ester. MS: m/z = 558.6 M+1

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-dimethylamino-pyrrolidin-1-yl)-4-oxo-butamide. MS: $m/z = 494.5$ $M+1$

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(2-oxa-5-aza-bicyclo[2.2.1]hept-5-yl)-4-oxo-butamide. MS: $m/z = 471$ $M+1$


N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(2,6-dimethyl-morpholin-4-yl)-4-oxo-butamide. MS: $m/z = 487$ $M+1$

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(2-methoxymethyl-morpholin-4-yl)-4-oxo-butamide. MS: $m/z = 503$ $M+1$.

EXAMPLE 10


N-(1-[(Benzyl-methyl-amino)-methyl]-1-cyano-3-phenyl-propyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide

(a) (Benzyl-methyl-amino)-acetic acid methyl ester




Methyl bromoacetate (10 g, 65.4 mmol, 1.0 equiv) was dissolved in 100 mL of CH_2Cl_2 which was then cooled to 0°C . *N*-Methylbenzylamine (7.92 g, 65.4 mmol, 1.0 equiv) was added dropwise over a 10 min period at which time Hunig's base (21 g, 163 mmol, 2.49 equiv) was added and the reaction was allowed to warm to room temperature and was stirred for 16 h. The reaction was diluted with 50 mL of saturated bicarbonate and the layers were separated. The organic layer was washed once more with 50 mL of sodium bicarbonate, dried over Na_2SO_4 , decanted and concentrated to a yellow oil which was used in the next step without further purification.

(b) (Benzyl-methyl-amino)-acetic acid hydrochloride salt.




(Benzyl-methyl-amino)-acetic acid methyl ester (12 g, 62.1 mmol, 1.0 equiv) was dissolved in 50 mL of MeOH to which was added a solution of LiOH hydrate (8.23 g, 196 mmol, 3.0 equiv) in 50 mL of water. The mixture was stirred at room temperature for 1.3 h at which time TLC (5% MeOH in CH₂Cl₂) indicated consumption of the starting material. The reaction solution was washed 1 x 50 mL of Et₂O and then the aqueous layer was acidified to pH 1 with concentrated HCl. The mixture was then evaporated to dryness under high vacuum. The resulting solid was triturated with EtOH (200 mL) and the mixture was filtered. The filtrate was then concentrated to dryness to yield a white crystalline solid that was used without further purification.

(c) 2-(Benzyl-methyl-amino)-*N*-methoxy-*N*-methy-acetamide.




(Benzyl-methyl-amino)-acetic acid hydrochloride salt (12 g, 56 mmol, 1.0 equiv) was mixed in DMF (200 mL) along with EDC (10.6 g, 56 mmol, 1.0 equiv) for 15 min. *N,O*-Dimethylhydroxylamine hydrochloride (6.52g, 67 mmol, 1.2 equiv) was added to the reaction followed by *N*-methylmorpholine (34 mL, 334 mmol, 6.0 equiv). The reaction was stirred for 16 h. The DMF was removed in vacuo and the residue was redissolved in 200 mL of EtOAc. The solution was washed once with 200 mL brine. The organic layer was dried over Na₂SO₄, decanted and concentrated to yield the desired product as a free-flowing oil (12.6 g, 80%) that was used in the next step without further purification; MS, *m/z* 223=*M*+1.

(d) 1-(Benzyl-methyl-amino)-4-phenyl-butan-2-one.



2-(Benzyl-methyl-amino)-*N*-methoxy-*N*-methyl-acetamide (2.0 g, 9.0 mmol, 1.0 equiv) was dissolved in 30 mL of dry THF under argon. The solution was cooled to -78°C and a 1 M solution of phenethyl magnesium chloride (12.6 mL, 12.6 mmol, 1.4 equiv) was added dropwise over a period of 1 min. The resulting mixture was stirred at -78°C for 3 h, warmed to 0°C for 20 min and re-cooled to -78°C at which time the reaction was quenched by the addition of 5 mL of saturated sodium bicarbonate. The reaction solution was diluted with 200 mL of EtOAc and 200 mL of water. The layers were separated and the organic layer was dried over Na_2SO_4 , decanted and concentrated to an oil (2.4 g, $\sim 100\%$) which was used without further purification; MS, m/z 268=M+1.


(e) 2-Amino-2-[(benzyl-methyl-amino)-methyl]-4-phenyl-butyronitrile.



1-(Benzyl-methyl-amino)-4-phenyl-butan-2-one (2.4 g, 9.0 mmol, 1.0 equiv) was mixed in 36 mL of 2.0 M NH_3 in MeOH. To the solution was added NaCN (0.484 g, 9.9 mmol, 1.1 equiv) and NH_4Cl (0.528 g, 9.9 mmol, 1.1 equiv). The resulting mixture was refluxed for 2 h at which time an addition 15 mL of 2.0 M NH_3 in MeOH was added and the mixture was refluxed another 4 h. Though TLC (5% MeOH in CH_2Cl_2) still showed the presence of the ketone, the reaction was cooled and filtered. The filtrate was concentrated and the crude product was purified by

flash chromatography on SiO₂ using 60% hexanes in CH₂Cl₂ then 1.5% MeOH to 3% MeOH was added. The desired product was isolated pure (0.561g, 21%) as a yellow oil that crystallized upon standing at -20 °C; MS, m/z 294=M+1.

(f) *N*-{1-[(benzyl-methyl-amino)-methyl]-1-cyano-3-phenyl-propyl}-2-cyclohexylmethyl-4-morpholin-4-yl-oxo-butamide.




2-Cyclohexylmethyl-4-morpholin-4-yl-oxo-butyanoic acid (0.097 g, 0.34 mmol, 1.0 equiv) was dissolved in 10 mL of CH₂Cl₂ followed by addition of EDC (0.065 g, 0.34 mmol, 1.0 equiv). The solution was stirred for 15 min at which time a solution of 2-amino-2-[(benzyl-methyl-amino)-methyl]-4-phenyl-butyronitrile (0.10 g, 0.34 mmol, 1.0 equiv) in 5 mL of CH₂Cl₂ was added followed by *N*-methylmorpholine (0.137 g, 1.36 mmol, 4.0 equiv). The resulting solution was stirred at room temperature for 72 h. The volatiles were removed *in vacuo* and the residue was redissolved in 100 mL of EtOAc and the solution was washed 2 x 25 mL brine. The organic layer was evaporated to dryness and the crude product was purified by semi-prep reverse-phase HPLC using 40% CH₃CN in water to 90% over a time of 25 min. The appropriate fractions were combined and concentrated to yield the desired product as a thick oil (0.047g, 25%); MS, m/z 559=M+1.

EXAMPLE 11


N-(4-Cyano-1,2-dimethyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide.

(a) 3-(Benzyl-methyl-amino)-butyric acid methyl ester.




Benzyl-methyl-amine (20 g, 165 mmol, 1.0 equiv) was added neat to methyl crotonate (19.8 g, 198 mmol, 1.2 equiv). The resulting solution was stirred at room temperature for 72 h. The excess crotonic ester was removed *in vacuo* to yield the desired product (40.3 g, ~100%) which was used without further purification; MS, m/z 222=M+1.

(b) 3-Methylamino-butyric acid methyl ester.




3-(Benzyl-methyl-amino)-butyric acid methyl ester (15 g, 67.8 mmol, 1.0 equiv) was placed in a Parr hydrogenation bottle and dissolved in 50 mL of MeOH. 20% Palladium hydroxide on carbon (0.5 g, 0.94 mmol, 0.014 equiv) was added and the mixture was shaken at 50 psi H_2 for 16 h. The reaction was judged as complete when the uptake of H_2 had stopped. The bottle was opened and 10 g of diatomaceous earth in 100 mL of MeOH was added. The mixture was filtered on a pad of diatomaceous earth which was then washed with 2 x 100 mL of MeOH. The filtrates were combined and concentrated *in vacuo* to yield the desired product as an oil that is somewhat volatile (7.6 g, 85%). The crude product was used without further purification; MS, m/z 132=M+1.

(c) 3-[(2-Methoxycarbonyl-ethyl)-methyl-amino]-butyric acid methyl ester.



3-Methylamino-butyric acid methyl ester (7.6 g, 58 mmol, 1.0 equiv) was added neat to methyl acrylate (7.5g, 87 mmol, 1.5 equiv). The resulting solution was refluxed for 16 h. The reaction was cooled and diluted with hexanes (200 mL) and an insoluble polymer separated out. The hexane solution was decanted and the polymer was washed 2 x 100 mL hexanes with vigorous stirring. The combined hexane solutions were then concentrated *in vacuo*. The crude product was purified by flash chromatography on SiO₂ using pure CH₂Cl₂ as an eluent. The pure product was isolated as a clear colorless oil (7.3g, 58%); MS, *m/z* 218=M+1.


(d) **1,2-Dimethyl-4-piperidone.**



A 1 M solution of TiCl₄ in CH₂Cl₂ (23 mL, 23 mmol, 1.0 equiv) was added to a flask under Ar and cooled to -15 °C with a MeOH/ice water bath. 3-[(2-Methoxycarbonyl-ethyl)-methylamino]-butyric acid methyl ester (5 g, 23 mmol, 1.0 equiv) was added dropwise over a 25 min period as solution in 75 mL of dry CH₂Cl₂ to give a dark red mixture that was difficult to stir with a magnetic stir bar. Stirring was continued an additional 1 h and then Et₃N (5.1g, 50.6 mmol, 2.2 equiv) was added dropwise over a 30 min period and then the reaction was stirred an additional 1.5 h at -15 °C. The reaction mixture was poured into 150 mL of brine and 150 mL of CH₂Cl₂ was added. After thorough mixing, the pH of the water was brought to 8-9 with Et₃N. The mixture was filtered and the gel-like solid was washed 3 x 100 mL CH₂Cl₂. The filtrate layers were separated and the aqueous layer was washed with 3 x 50 mL CH₂Cl₂. All of the organic layers were combined and concentrated to a thick red oil. The residue was taken up in 150 mL of concentrated HCl and the solution was refluxed 4 h. The cooled reaction solution was evaporated to dryness and the residue was dissolved in 200 mL of saturated sodium bicarbonate. The product ketone was extracted with 2 x 100 mL of EtOAc. The organic layers were combined and dried over Na₂SO₄. The product was purified by flash chromatography on SiO₂


using CH_2Cl_2 to 4% MeOH in CH_2Cl_2 as eluent. The product was isolated as an orange oil (1.23 g, 42%); MS, m/z 128= $M+1$.

(e) 4-Amino-4-cyano-1,2-dimethyl-piperidine.



1,2-Dimethyl-4-piperidone (1.23 g, 9.67 mmol, 1.0 equiv) was dissolved in 39 mL of 2 M NH_3 in MeOH (8 equiv NH_3). To this solution was added NaCN (0.52 g, 10.6 mmol, 1.1 equiv) and NH_4Cl (0.57 g, 10.6 mmol, 1.1 equiv). The resulting mixture was refluxed for 2 h at which time an additional 39 mL of 2 M NH_3 in MeOH was added followed by an additional 2 h of reflux. The reaction was cooled and filtered. The filtrate was concentrated and taken up in 100 mL of CH_2Cl_2 giving more salt precipitate which was removed by a second filtration. The filtrate was then concentrated to a thick orange oil (1.32 g, 89%). ^1H NMR showed a 3 to 1 mixture of diastereomers of unknown configuration. The crude product was used without further purification; MS, m/z 154= $M+1$.

(f) *N*-(4-Cyano-1,2-dimethyl-piperidin-4-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide.



2-Cyclohexylmethyl-4-morpholin-4-yl-oxo-butanoic acid (0.3 g, 1.06 mmol, 1.0 equiv) was mixed with EDC (0.2 g, 1.06 mmol, 1.0 equiv) in 15 mL of DMF. A solution of 4-amino-4-cyano-1,2-dimethyl-piperidine (0.16 g, 1.0 mmol, 0.95 equiv) in 5 mL of DMF was added

followed by addition of *N*-methylmorpholine (0.43 g, 4.24 mmol, 4.0 equiv). The resulting solution was stirred for 16 h. The reaction was diluted with 50 mL of saturated sodium bicarbonate and the product was extracted with 3 x 50 mL EtOAc. The organic layers were combined and concentrated *in vacuo*. The crude product was purified by semi-prep reverse-phase HPLC using 20% CH₃CN in water to 61% CH₃CN in water over a gradient time of 17 min. The product was isolated in two peaks as an undefined mixture of diastereomers about the piperidine ring; MS, *m/z* 419=*M*+1 for both HPLC peaks.

EXAMPLE 12

(4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butylamide

(a) 3-(4-Methyl-cyclohexyl)-acrylic acid ethyl ester

Lithium aluminium hydride (560 mg, 60 mmol) was suspended in 20 mL of dry THF. To this suspension at -78 °C under nitrogen was added a solution of *trans*-4-methyl-1-cyclohexanecarboxylic acid (2.13 g, 15 mmol) in 20 mL of dry THF. This reaction mixture was stirred at -78 °C for 2 h and then warmed up to room temperature in 2 h. The reaction was cooled to 0 °C, and quenched with a saturated solution of potassium sodium tartrate and extracted repeatedly with diethyl ether. The combined organic phases were dried over magnesium sulfate and evaporated under reduced pressure to give a clear oil. Yield 100%.

To a solution of oxalyl chloride (1.76 mL, 20 mmol) in 60 mL of dry dichloromethane at -78 °C under nitrogen was added DMSO (2.84 mL, 40 mmol). After the gas evolution stopped, the above oil (1.28 g, 10 mmol) dissolved in 10 mL of dichloromethane was added. The reaction mixture was stirred at -78 °C for 1 h and triethylamine (7 mL, 50 mmol) was added. The reaction was then allowed to warm up to room temperature in 2 h and was quenched with water. The aqueous phase was extracted repeatedly with dichloromethane. The combined organic phases were washed with brine and dried over magnesium sulfate. The

solvent was removed under reduced pressure to give a yellow oil, which was used in the next step without further purification. Yield: 1.06 g, 84%.

The above oil (1.06 g, 8.41 mmol) and (carbethoxymethylene)triphenylphosphorane (2.93 g, 8.41 mmol) were heated under reflux in 30 mL of dry toluene under nitrogen for 2 h. The solvent was removed under reduced pressure and the residue was purified by silica gel flash chromatography eluting with 5% diethyl ether in petroleum ether to give a clear oil. Yield: 1.45 g, 88%.

(b) 3-(4-Methyl-cyclohexyl)-propionic acid.

The above oil (1.45 g, 7.39 mmol) was dissolved in 20 mL of EtOH and hydrogenated in a Parr apparatus under 50 psi of H₂ in the presence of 10 % palladium on activated charcoal for 2 h. The solid was removed by filtration and washed with EtOH. The filtrate was evaporated under reduced pressure to give the ethyl ester as a clear oil. Yield: 1.37 g, 94%.

The above ester (1.37 g, 6.98 mmol) was dissolved in 20 mL of THF. A 2 M lithium hydroxide solution (50 mL) was added. This reaction mixture was stirred at room temperature for 16 h. The solution was cooled to 0 °C, and acidified with 6 N HCl to pH 2. The resulting white precipitate was collected by filtration and washed with water then dried under vacuum giving the desired acid. Yield: 1.08 g, 91%.

(c) (S)-4-Isopropyl-3-[3-(4-methyl-cyclohexyl)-propionyl]-oxazolidin-2-one.

The acid (1.02, 6 mmol) was dissolved in 20 mL of dry dichloromethane and 1 drop of DMF. To this solution at 0 °C was added oxalyl chloride (1.57 mL, 18 mmol). The reaction mixture was stirred at room temperature for 2 h. The solvent was removed under reduced pressure and the resulting acid chloride was used in the next reaction without further purification.

(*S*)-(-)-4-Isopropyl-2-oxazolidinone (0.775 g, 6 mmol) was dissolved in 20 mL of dry THF. To this solution at -78 °C, under nitrogen was added *n*-butyl lithium (2.5 M, 2.4 mL, 6 mmol). The reaction mixture was stirred at -78 °C for 30 min and then at room temperature for 30 min. To this solution at -78 °C was added the acid chloride dissolved in 20 mL of THF. The reaction was maintained at -78 °C for 1 h then warmed up to room temperature in 2 h and quenched with 1M potassium carbonate solution. The aqueous phase was extracted repeatedly with diethyl ether. The combined organic phases were washed with brine and dried over magnesium sulfate. The solvent was removed under reduced pressure. The residue was purified by silica gel flash chromatography, eluted with 5% diethyl ether in petroleum ether to give a clear oil. Yield: 1.39 g, 82 %.

(d) (2*R*,4'*S*)-1-(4-Isopropyl-2-oxo-oxazolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-butane-1,4-dione.

The above compound (0.703 g, 2.5 mmol) was dissolved in 20 mL of dry THF. To this solution at -78 °C and under nitrogen was added sodium bis(trimethylsilyl)amide (0.6 M, 4.2 mL, 2.5 mmol). The reaction mixture was stirred at -78 °C for 1 h. 2-Bromo-1-morpholin-4-yl-ethanone (W.J. Thompson, et al., *J. Med. Chem.*, 1992 35 1685-1701) (0.52 g, 2.5 mmol) dissolved in 10 mL of dry THF was added to the above reaction mixture. This solution was maintained at -78 °C for 1 h, warmed up to room temperature in 2 h and quenched with saturated ammonium chloride solution. The aqueous phase was extracted repeatedly with dichloromethane. The combined organic phases were washed with brine and dried over magnesium sulfate. The solvent was removed under reduced pressure. The residue was purified by silica gel flash chromatography, eluting with 1% MeOH in dichloromethane to give a white solid. Yield: 0.82 g, 80 %.

(e) (*R*)-2-(4-Methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyric acid.

The above solid (0.82 g, 1.96 mmol) was dissolved in 30 mL of THF and 10 mL of water. To this solution, at 0 °C was added 30 mL of 30% hydrogen peroxide. The reaction mixture was stirred at 0 °C for 2 h and then quenched with sodium sulfite. This solution was acidified with 6 N HCl to pH 2 and then extracted repeatedly with dichloromethane. The

combined organic phases were washed with brine and dried over magnesium sulfate. The solvent was removed under reduced pressure. The residue was purified by silica gel flash chromatography, eluted with 2% MeOH in dichloromethane to give a white solid. Yield: 0.31 g, 53 %.

(f) (4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide.

The title compound was prepared using the above intermediate following the procedure described in Example 6, C₂₇H₄₄N₄O₃, ESMS: 473 (M+1)

Additional compounds of the present invention prepared by methods analogous to those above include:

(4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(1-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide: C₂₇H₄₄N₄O₃, ESMS: 473 (M+1)

EXAMPLE 13

***N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(*trans*-4-phenyl-cyclohexylmethyl)-butyramide.**

(a) 4-Phenyl-cyclohexanecarboxylic acid.

To a mixture of AlCl₃ (13 g, 97.5 mmol, 1.0 equiv) in benzene (100 mL) kept at 35 °C, was added 1-cyclohexenecarboxylic acid (5 g, 39.7 mmol, 0.41 equiv) dropwise. After addition was complete, the reaction solution was heated to 70 °C for 1 h and 45 °C overnight. The reaction was cooled, and 200 mL of 1 N HCl was slowly added. The product was extracted with EtOAc. The organic layer was extracted with 1 N NaOH and the aqueous layer was acidified with concentrated HCl to produce a white precipitate that was collected by filtration, washed with water and dried to give the titled compound as a white solid (3.1 g, 38%) which was used without further purification.

(b) 4-Phenyl-cyclohexylmethanol.

A suspension of LiAlH_4 (0.56 g, 14.8 mmol, 2.0 equiv) was prepared under argon in 50 mL of anhydrous THF (50 mL) kept at 0 °C. A solution of 4-phenyl-cyclohexanecarboxylic acid (1.5 g, 7.34 mmol, 1.0 equiv) in 50 mL of THF was added dropwise over a 20 min period. The reaction mixture was stirred at ambient temperature for 1.5 h and then cooled to 0 °C. The excess hydride reagent was carefully quenched under argon by dropwise addition of 1 N HCl. After quenching, the reaction mixture was extracted with EtOAc. The organic layer was dried over MgSO_4 , filtered and concentrated to give the title product as a colorless oil (1.9 g crude) which was used without further purification.

(c) 4-Phenyl-cyclohexylmethyl-*p*-toluene sulfonate.

4-Phenyl-cyclohexylmethanol (2.5 g, 13.1 mmol, 1.0 equiv) was dissolved in chloroform (50 mL) and pyridine (2.1 g, 26.3 mmol, 2.0 equiv) was added and the mixture was cooled to 0 °C. A solution of tosyl chloride (3.75 g, 19.7 mmol, 1.5 equiv) in chloroform (30 mL) was added and the resulting solution was stirred at ambient temperature for 30 min. The reaction was diluted with EtOAc (150 mL) and washed 2 x 100 mL 1N HCl. The organic layer was then washed with 1 x 100 mL saturated sodium bicarbonate, dried over MgSO_4 , filtered and concentrated to yield a solid which was used immediately in the next step without further purification.

(d) (4-Phenyl-cyclohexylmethyl)-malonic acid diethyl ester.

Diethyl malonate (3.1 g, 19.7 mmol, 1.5 equiv) was added dropwise to a solution of sodium (1.25 g, 54.0 mmol, 4.0 equiv) in EtOH (50 mL). 4-Phenyl-cyclohexylmethyl-*p*-toluene sulfonate (~13.1 mmol, ~1.0 equiv, the crude product from the previous step) was added as a solution in 50 mL of EtOH. The resulting mixture was refluxed for 8 h. The reaction mixture was cooled and evaporated to dryness and then dissolved in water (150 mL) and the crude product extracted with 3 x 100 mL EtOAc. The organic layer was dried over MgSO_4 , filtered and concentrated. The crude product was purified by flash

chromatography on SiO₂ using 10% EtOAc in hexanes to give the desired product as a colorless oil (2.79 g).

(e) 3-(4-Phenyl-cyclohexylmethyl)-propanoic acid.

(4-Phenyl-cyclohexylmethyl)-malonic acid diethyl ester (2.79 g, 8.4 mmol, 1.0 equiv) was added to a solution of potassium hydroxide (2.35 g, 41 mmol, 5 equiv) in 20 mL of water and 20 mL of MeOH. The resulting mixture was refluxed for 0.5 h and then evaporated to dryness and then dissolved in 100 mL of water. The solution was washed 3 x 100 mL with hexane and then brought to pH <1 with concentrated HCl. The product was extracted 3 x 100 mL EtOAc. The organic layers were combined and concentrated to dryness. The crude product was then heated for 10 min at 180° C to decarboxylate the diacid and yield the monoacid (0.9 g) which was used without further purification.

(f) *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(*trans*-4-phenyl-cyclohexylmethyl)-butyramide;

The title compound was prepared using the intermediate from step e above and the procedures reported in Example 12, MS, *m/z* 535=M+1

EXAMPLE 14

2-Bicyclo[4.1.0]hept-7-ylmethyl-N-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butamide.

(a) 2-Bicyclo[4.1.0]hept-7-yl-carboxylic acid ethyl ester.

A mixture of cupric triflate (0.1 g, 0.28 mmol, 0.0016 equiv) was prepared in cyclohexene (100 mL). A 25 mL of a solution of ethyl diazoacetate (20 g, 175 mmol, 1.0 equiv) in cyclohexene (100 mL) was added to initiate the reaction followed by dropwise addition of the remainder to

maintain a gentle exothermic reaction. The reaction was then stirred 18 h at room temperature. 200 mg of additional cupric triflate was added to decompose any unreacted diazo acetate and then the mixture was concentrated to dryness. The residue was distilled on a water bath to yield the desired product as a colorless oil (15.7 g).

(b) 2-Bicyclo[4.1.0]hept-7-ylmethyl-N-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butyramide.

2-Bicyclo[4.1.0]hept-7-ylmethyl-N-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butyramide was prepared using the procedures outlined in Example 13 and the intermediate generated in step a above; MS, m/z 471=M+1.

Additional compounds of the present invention prepared by methods analogous to those above include:

2-Bicyclo[4.1.0]hept-7-ylmethyl-N-(4-cyano-1-methyl-piperidin-4-yl)-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 417=M+1.


N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-butyramide; MS, m/z 507=M+1 .

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-indan-2-ylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 493=M+1.

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclopentylmethyl-4-morpholin-4-yl-4-oxo-butyramide; MS, m/z 445=M+1.

EXAMPLE 15

[2-Cyano-2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-ethyl]-carbamic acid tert-butyl ester.




- (a) 3-tert-Butoxycarbonylamino-2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-propionic acid methyl ester was prepared by the procedure described in Example 1 from (R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyric acid (0.25 g, 0.88 mmol), EDC (0.22 g, 1.15 mmol), 1-hydroxybenzotriazole (0.155 g, 0.1.15 mmol) and the HCl salt of N-Beta-Boc-Alpha-Beta-Diaminopropionic acid methyl ester (0.25 g, 0.98 mmol) to yield a white solid (0.36 g, 84.5%) after purification by flash column chromatography. MS: m/z 484 = $M+1$
- (b) The above product (0.36 g, 0.745 mmol) was dissolved in 25 mL of 2M ammonia in MeOH. The reaction was cooled to 5°C and saturated with ammonia. The reaction was stoppered and left at room temperature overnight. The volatiles were removed to give the corresponding amide [2-Carbamoyl-2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-ethyl]-carbamic acid tert-butyl ester (0.34g, 97.5%). MS: m/z 469 = $M+1$
- (c) The amide from the step above (0.050 g, 0.107 mmol) and cyanuric chloride (0.020, 0.108 mmol) were reacted as described by the procedure in Example 1. The residue was purified by chromatography (SiO_2 , 5% MeOH in CH_2Cl_2) to give the title compound as a white solid

(0.02, 48%); ^1H NMR (CDCl_3) 6.83-6.81(1H, d), 5.49(1H, m), 4.93(1H, s), 3.68-3.44(8H, m), 2.79-2.73(2H, m), 2.32(1H, d), 1.7-1.59(8H, m), 1.44(9H, s), 1.25-1.11(4H, m), 0.89-0.86(3H, m). MS: m/z 451 = $M+1$

EXAMPLE 16

N-{Cyano-[(cyclohexyl-ethyl-amino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide.



- (a) To [2-Carbamoyl-2-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butrylamino)-ethyl]-carbamic acid tert-butyl ester (0.15g, 0.32 mmol) in 5 mL methylene chloride at room temperature was added 0.5 mL of TFA. The reaction was stirred for 1h and the volatiles removed. The residue was used for the next step without further purification. MS: m/z = 369 $M+1$.
- (b) To the above N-(2-Amino-1-carbamoyl-ethyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide TFA salt (0.154g, 0.32 mmol) in 10 mL THF was added cyclohexanone (0.62g, 0.64 mmol). The reaction was cooled to 0°C and sodium triacetoxymethylborohydride was added

(0.176 g, 0.83 mmol) and the reaction was allowed to warm up to room temperature overnight. The reaction mixture was cooled in an ice bath and acetaldehyde (0.1 mL, excess) was added. The ice bath was removed and the reaction was stirred for 2h. The reaction was quenched with saturated sodium bicarbonate solution, extracted with EtOAc and dried over anhydrous sodium sulfate. The volatiles were removed and the crude product was purified by flash column chromatography (SiO₂, 5% MeOH in CH₂Cl₂) to give the desired product (0.1g, 65.5%). MS: m/z = 479 M+1.


- (c) The amide from the step above (0.070 g, 0.146 mmol) and cyanuric chloride (0.054 g, 0.292 mmol) were reacted as described by the procedure in Example 1. The residue was purified by chromatography (SiO₂, 5% MeOH in CH₂Cl₂) to give the title compound as a white solid (0.035g, 51.9%); ¹H NMR (CDCl₃) , 4.64-4.59(1H, m), 3.66-3.62(6H, m), 3.56(2H, m), 3.45(2H, m), 2.86-2.7(5H, m), 2.62(2H, m), 2.5(1H, m), 2.3(1H, m), 1.78(6H, m), 1.6(5H, m), 1.26(6H, m), 1.1(3H, t), 0.85(3H, m). MS: m/z = 461 M+1

The following compound was synthesized using the above procedure:

N-{Cyano-[(dibenzylamino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, MS: m/z = 531 M+1

EXAMPLE 17


N-([(Benzyl-ethyl-amino)-methyl]-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.



- (a) To 1N HCl/MeOH (50 mL) at 0°C was added (S)-2-Amino-3-benzylamino-propanoic acid (0.5g, 2.57 mmol). Reaction was stirred at room temperature overnight. Volatiles were removed to give the product 2-Amino-3-benzylamino-propanoic acid methyl ester dihydrochloride salt as a white solid (0.7g, 97.5%). MS: $m/z = 209 M+1$
- (b) The title compound was synthesized according to the scheme using the procedures in Example 2. The diastereomers were separated via flash column chromatography to give N-[[(Benzyl-ethyl-amino)-methyl]-cyano-methyl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide, MS: $m/z = 469 M+1$

EXAMPLE 18

N-(Cyano-piperidin-1-ylmethyl-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide.



- (a) To sodium carbonate (1.13 g, 10.67 mmol) in water (10 mL) and THF (15 mL) at 0°C was added L-serine amide hydrochloride (1.0 g, 7.1 mmol). The reaction was stirred for 5 min and then benzyl chloroformate was added (1.2g, 7.1 mmol). The reaction was allowed to warm up to room temperature and stirred for 4 h. Volatiles were removed and the residue was extracted with methylene chloride/EtOAc (2:1). The organic fraction was dried over anhydrous sodium sulfate and evaporated to give the CBz protected serine amide as a white solid (1.0g, 59%). NMR and MS were consistent with the desired product.
- (b) To the above CBz protected serine amide (0.5 g, 2.1 mmol) in dry pyridine (3 mL) at 0°C was added methanesulfonyl chloride (0.2 mL, 2.52 mmol) dropwise. After addition was complete, the reaction was stirred for 2 h at 0°C. It was poured over 10 mL of cold aqueous 1N HCl and extracted with EtOAc (2x25 mL). The combined organic fractions were washed with saturated sodium bicarbonate, brine and dried over anhydrous sodium sulfate. Removal of volatiles afforded the desired product as a white solid (0.6 g, 90.5%). NMR and MS were consistent with the desired product.
- (c) The above mesylate (0.4g, 1.26 mmol) was dissolved in MeOH (8 mL) at 0°C and to this was added sodium carbonate (0.13 g, 1.26 mmol) and piperidine (0.11 g, 1.26 mmol). The reaction was stirred at 0°C for 6 h. Solvent was removed and the residue was extracted with EtOAc. The solid was filtered off, filtrate evaporated and purified by flash column chromatography (Silica gel, 5%CH₃OH/CH₂Cl₂). The desired piperidine serine amide was

obtained as a white solid (0.3 g, 77%). NMR and MS consistent with desired product. MS: $m/z = 306 M+1$.

(d) The above piperidine compound (0.09 g, 0.29 mmol) was dissolved in MeOH (15 mL) and subjected to transfer hydrogenation using cyclohexene (3 mL) and palladium hydroxide on carbon (0.020 g). The reaction was heated at 70°C for 45 min., cooled and filtered over diatomaceous earth. Removal of solvent gave the free amine (0.04 g, 79.3%). NMR and MS consistent with desired product. MS: $m/z = 172 M+1$.

(e) The above amine was further coupled and then dehydrated using conditions described in Example 2, giving the target compound: N-(Cyano-piperidin-1-ylmethyl-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide, 1H NMR ($CDCl_3$) δ 4.75(1H, m), 3.67(5H, m), 3.55(2H, m), 3.44(1H, m), 2.7(3H, m), 2.57(2H, m), 2.45(2H, m), 2.31(1H, m), 1.7(6H, m), 1.4(3H, m), 1.2(8H, m), 0.89(3H, m). MS: $m/z = 419 M+1$.

EXAMPLE 19

N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide.


N-(1-Benzyl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide was prepared via a slight modification of Example 6. A 3-tert-butoxycarbonyl amino pyrrolidine was coupled to (R)-2-cyclohexyl methyl succinic acid 1-methyl ester. The Boc protecting group was removed with HCl/dioxane and the resulting amino pyrrolidine was alkylated with methanesulfonyl chloride. This intermediate was then carried through the standard synthetic sequence outlined in Example 6 to give the title compound; MS: $m/z = 544.4 M+1$.

The following compound was synthesized using the above procedure:

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-(3-methanesulfonylamino-pyrrolidin-1-yl)-4-oxo-butyramide. MS: $m/z = 536.4 M+1$.

EXAMPLE 20

N-[(3R)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide.



(a) N-Benzyloxycarbonyl-(R)-3-hydroxypyrrolidine.

To a stirred mixture of (R)-3-hydroxypyrrolidine (4.35 g, 50 mmol) in THF (50 mL) and aqueous sodium carbonate (10.6 g, 100 mmol in 50 mL water) cooled to 0°C was added a solution of benzyl chloroformate in THF (60 mmol, 8.6 mL in 10 mL THF) over a period of 5-10 min. After stirring at 0°C for 2 h, the reaction mixture was diluted with 200 mL water and extracted with methylene chloride (3X100 mL). The combined extracts were washed with 1N sulfuric acid (3X75 mL) followed by brine solution (3X50 mL). This extract was

then dried over anhydrous sodium sulfate and the solvent evaporated to give a light yellow oil (10.5 g, 95% yield) which was used for the next reaction without any purification.

(b) (3R)-3-Cyano-3-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-pyrrolidine-1-carboxylic acid benzyl ester.

The title compound was prepared using the procedures outlined in Example 6 and the intermediate generated from step a. above.

(c) Hydrogenolysis of (3R)-3-cyano-3-(2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyrylamino)-pyrrolidine-1-carboxylic acid benzyl ester.

A solution of the CBZ-compd., (1.8 g, 3.53 mmol) in EtOH (75 mL) was stirred under an atmosphere of hydrogen gas along with 10% palladium-carbon (0.3 g). After 1h catalyst was filtered through a diatomaceous earth and the solvent evaporated to give a thick colorless oil (1.26 g, 95%) and was used without purification.

(d) Reductive amination

To a stirred solution of the amine (0.3 mmol, 113 mg) and the 3,3-dimethylcyclohexanone (1 mmol, 126 mg) in 1,2-dichloroethane (1 mL) was added sodium triacetoxyborohydride (0.3 mmol, 66 mg). After 2 h the reaction mixture was diluted with 10 mL methylene chloride, washed with water (3x5 mL) and the solvent was evaporated from the organic layer to give a colorless oil which was taken up in 50% aqueous acetonitrile and purified by reverse phase hplc. The later eluting peak was characterized to be N-[(3R)-3-cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide; (MH+ 487, ¹H NMR (DMSO-d₆) 0.5-1.7 (27H, m), 1.9-2.6 (8H, m), 2.6, 2.85 (1H, dd), 3 (1H, d), 3.15-3.34 (8H, m), 8.45 (1H, s).

The following compounds were synthesized using the above procedure:

N-[(3S)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[1-trans-(4-tert-Butyl-cyclohexyl)-(3R)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[1-cis-(4-tert-Butyl-cyclohexyl)-(3S)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[1-cis-(4-tert-Butyl-cyclohexyl)-(3R)-3--cyano-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3S)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

N-[(3R)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide


N-[(3S)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3S)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butylamide

N-[(3R)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide

EXAMPLE 21

N-(2-Benzoyloxymethyl-1-cyano-cyclopropyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butamide

To a stirred mixture of NaH (60% dispersion in mineral oil, 35.9 g, 149.5 mmol) in DME (100 mL) at 0 °C, was added a solution of 4-benzyloxymethyl-[1,3,2]-dioxathiolane-2,2-dioxide (K. Burgess et al., *J. Org. Chem.* 1993, 58, 3767-3768) (17.4 g, 71.2 mmol) and (benzhydrylidene-amino)acetic acid ethyl ester (19.0 g, 71.2 mmol) in DME (100 mL). The mixture was warmed to 55 °C (30 min later, the mixture turned to solid and thus 200 mL of DME was added) and stirred for 4 h. It was allowed to cool to room temperature, slowly poured onto ice in saturated aqueous ammonium chloride solution, extracted with dichloromethane, washed with brine, dried (sodium sulfate) and concentrated to give crude ester product.

A mixture of the above ester and *p*-toluenesulfonic acid monohydrate (13.5 g, 71.2 mmol) in MeOH (170 mL) and water (25 mL) was stirred at room temperature for 1 h. It was diluted with water, extracted with dichloromethane, washed with brine, dried (sodium sulfate) and purified by flash chromatography on silica gel (4 hexane/1 EtOAc) to give the desired amino ester (11.0 g, 61.9% in two steps).

Typical peptide coupling of the above amino ester with (*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butanoic acid gave the desired coupled product in 50% yield.

A mixture of the above intermediate (13.1 g, 25.5 mmol) and lithium hydroxide (1.2 g, 2 equiv.) in MeOH (100 mL) and water (30 mL) was stirred at room temperature overnight. It was diluted with 1N HCl, extracted with dichloromethane, washed with brine, dried (sodium sulfate) and concentrated to dryness to give the desired carboxylic acid (10.6 g, 86%).


To a stirred solution of the above acid (11.3 g, 23.3 mmol) in THF (50 mL) at -10°C was added triethylamine (3.9 mL, 1.2 equiv.) and ethyl chloroformate (2.7 mL, 1.2 equiv.). 20 Min later, ammonia gas was bubbled through the reaction for 4 h at -20°C . The reaction was allowed to warm to room temperature and stirred overnight. It was concentrated and purified by flash chromatography on silica gel (dichloromethane/MeOH = 10/1) to give the desired amide (11.3 g, 100%).

To a stirred solution of the above amide (178 mg, 0.36 mmol) and triethylamine (150 μL , 3 equiv.) in dichloromethane (10 mL) at 0°C was added TiCl_4 . The mixture was allowed to warm to room temperature and stirred for 20 min. It was washed with brine, dried (sodium sulfate) and purified by chromatography on silica gel (hexane/ethyl acetate = 1/1) to give the title compound (30 mg, 17.8%).

EXAMPLE 22

*N*¹-(1-Cyano-2-dimethylaminomethyl-cyclopropyl)-2-cyclohexylmethyl-*N*⁴-ethyl-*N*⁴-(2-methoxy-ethyl)-succinamide

5



A mixture of the intermediate amide from Example 21 (10 g, 20.6 mmol) and Pd/C (10%, 3 g) in MeOH (150 mL) was shaken under H₂ in a Parr apparatus overnight and filtered.
 10 The filtrate was concentrated to dryness to give the desired alcohol (7.8 g, 95.8%).

To a stirred solution of oxalyl chloride (2 M in dichloromethane, 7.6 mL, 15.2 mmol) in dichloromethane (30 mL) at -78 °C was added DMSO (2.2 mL, 31 mmol) dropwise. A solution of the above alcohol (6 g, 15.1 mmol) in dichloromethane (30 mL) was added. The mixture was allowed to warm to -40 °C in 1 h before triethylamine (8.5 mL, 61 mmol) was added. The mixture was then warmed to room temperature, washed with water and brine, dried (sodium sulfate) and concentrated to dryness to give the desired aldehyde (5.5 g, 92%).
 15


A mixture of the aldehyde from above (1 g, 2.5 mmol), *N*-benzylmethylamine (0.33 mL, 2.5 mmol) and sodium triacetoxyborohydride (0.81 g, 1.5 equiv.) in dichloromethane (10 mL) was stirred at room temperature overnight. The reaction
 20

mixture was washed with water, saturated aqueous ammonium chloride and brine, dried (sodium sulfate) and purified by flash chromatography on silica gel (dichloromethane/MeOH = 10/1) to give the desired benzyl methylamine (0.84 g, 66.8%).

A solution of the above amine (478 mg, 0.95 mmol) and cyanuric chloride (353 mg, 2 equiv.) in DMF (4 mL) was stirred at room temperature for 1 h. The reaction mixture was diluted with water, extracted with dichloromethane, washed with brine, dried (sodium sulfate) and purified by flash chromatography on silica gel (dichloromethane/MeOH = 20/1) to give the title compound (180 mg, 37.4%).

EXAMPLE 23

N-(1-Cyano-2-hydroxymethyl-cyclopropyl)-2-cyclohexylmethyl-4-morpholin-4-yl-2-oxo-butamide



To a stirred solution of the intermediate alcohol from Example 22 (1.1 g, 2.7 mmol) and pyridine (0.44 mL, 5.2 mmol) in dichloromethane (15 mL) at 0 °C was added Ti_2O (0.67 mL, 4 mmol) slowly. The mixture was allowed to warm to room temperature and stirred for 1 h. The reaction mixture was washed with brine, dried (sodium sulfate) and purified by chromatography on silica gel (dichloromethane/MeOH = 20/1) to give the title compound (0.35 g, 34.6%).

METHODS OF THERAPEUTIC USE

The compounds of the invention are useful in inhibiting the activity of cathepsin S, K, F,
5 L and cathepsin B. In doing so, these compounds are useful in blocking disease
processes mediated by these cysteine proteases.

Compounds of this invention effectively block degradation of the invariant chain to CLIP
by cathepsin S, and thus inhibit antigen presentation and antigen-specific immune
10 responses. Control of antigen specific immune responses is an attractive means for
treating autoimmune diseases and other undesirable T-cell mediated immune responses.
Thus, there is provided methods of treatment using the compounds of this invention for
such conditions. These encompass autoimmune diseases and other diseases involving
inappropriate antigen specific immune responses including, but not limited to,
15 rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, ulcerative colitis,
multiple sclerosis, Guillain-Barre syndrome, psoriasis, Grave's disease, myasthenia
gravis, scleroderma, glomerulonephritis, atopic dermatitis, insulin-dependent diabetes
mellitus and asthma. The compounds of the invention can also be used to treat other
disorders associated with extracellular proteolysis such as Alzheimer's disease and
20 atherosclerosis. The compounds of the invention can also be used to treat other disorders
associated with inappropriate autoimmune responses, T-cell mediated immune responses,
or extracellular proteolysis mediated by cathepsin S, unrelated to those listed above or
discussed in the Background of the Invention. Therefore, the invention also provides
methods of modulating an autoimmune disease comprising administering to a patient in
25 need of such treatment a pharmaceutically effect amount of a compound according to the
invention.

Compounds of the invention also include inhibition of cathepsin K. In doing so, they
may block inappropriate degradation of bone collagen and other bone matrix proteases.
30 Thus, there is provided a method for treating diseases where these processes play a role

such as osteoporosis. Inhibition of cathepsins F, L and B are also within the scope of the invention due to similarity of the active sites in cysteine proteases as described above.

For therapeutic use, the compounds of the invention may be administered in any
5 conventional dosage form in any conventional manner. Routes of administration include, but are not limited to, intravenously, intramuscularly, subcutaneously, intrasynovially, by infusion, sublingually, transdermally, orally, topically or by inhalation. The preferred modes of administration are oral and intravenous.

10 The compounds of this invention may be administered alone or in combination with adjuvants that enhance stability of the inhibitors, facilitate administration of pharmaceutical compositions containing them in certain embodiments, provide increased dissolution or dispersion, increase inhibitory activity, provide adjunct therapy, and the like, including other active ingredients. Advantageously, such combination therapies
15 utilize lower dosages of the conventional therapeutics, thus avoiding possible toxicity and adverse side effects incurred when those agents are used as monotherapies. Compounds of the invention may be physically combined with the conventional therapeutics or other adjuvants into a single pharmaceutical composition. Advantageously, the compounds may then be administered together in a single dosage form. In some embodiments, the
20 pharmaceutical compositions comprising such combinations of compounds contain at least about 15%, but more preferably at least about 20%, of a compound of formulas (I) or (Ia) (w/w) or a combination thereof. Alternatively, the compounds may be administered separately (either serially or in parallel). Separate dosing allows for greater flexibility in the dosing regime.

25

As mentioned above, dosage forms of the compounds of this invention include pharmaceutically acceptable carriers and adjuvants known to those of ordinary skill in the art. These carriers and adjuvants include, for example, ion exchangers, alumina,
aluminum stearate, lecithin, serum proteins, buffer substances, water, salts or electrolytes
30 and cellulose-based substances. Preferred dosage forms include, tablet, capsule, caplet, liquid, solution, suspension, emulsion, lozenges, syrup, reconstitutable powder, granule,

suppository and transdermal patch. Methods for preparing such dosage forms are known (see, for example, H.C. Ansel and N.G. Popovich, *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 5th ed., Lea and Febiger (1990)). Dosage levels and requirements are well-recognized in the art and may be selected by those of ordinary skill in the art from available methods and techniques suitable for a particular patient. In some embodiments, dosage levels range from about 10-1000 mg/dose for a 70 kg patient. Although one dose per day may be sufficient, up to 5 doses per day may be given. For oral doses, up to 2000 mg/day may be required. As the skilled artisan will appreciate, lower or higher doses may be required depending on particular factors. For instance, specific dosage and treatment regimens will depend on factors such as the patient's general health profile, the severity and course of the patient's disorder or disposition thereto, and the judgment of the treating physician.

ASSESSMENT OF BIOLOGICAL PROPERTIES

Expression and Purification of recombinant human Cathepsin S

Cloning of human cathepsin S:

U937 RNA was subjected to reverse transcriptase / polymerase chain reaction with primer A (5'cacaatgaaacggctggttg 3') and primer B (5'ctagatttctgggtaagaggg 3') designed to specifically amplify the cathepsin S cDNA. The resulting 900 bp DNA fragment was subcloned into pGEM-T (Promega) and sequenced to confirm its identity. This construct was used for all subsequent manipulations. This procedure is typical for cloning of known genes and is established in its field.

Human Pre-Pro-Cat S was removed from pGem-T vector (Promega, 2800 Woods Hollow Rd, Madison, WI 53711) by digestion with restriction enzyme SacII, followed by treatment with T4 DNA polymerase to generate a blunt end, and a second restriction enzyme digest with SalI. It was subcloned into pFastBac1 donor plasmid (GibcoBRL, 8717 Grovemont Cr., Gaithersburg, MD 20884) which had been cut with restriction enzyme BamHI and blunt-ended and then cut with restriction enzyme SalI. The ligation

mixture was used to transform DH5a competent cells (GibcoBRL) and plated on LB plates containing 100ug/ml ampicillin. Colonies were grown in overnight cultures of LB media containing 50ug/ml Ampicillin, plasmid DNA isolated and correct insert confirmed by restriction enzyme digestion. Recombinant pFastBac donor plasmid was transformed into DH10Bac competent cells (GibcoBRL). Large white colonies were picked from LB plates containing 50ug/ml kanamycin, 7ug/ml gentamicin, 10ug/ml tetracycline, 100ug/ml Bluo-gal, and 40ug/ml IPTG. DNA was isolated and used to transfect Sf9 insect cells using CellFECTIN reagent (GibcoBRL). Cells and supernatant were harvested after 72 hours. Viral supernatant was passaged twice and presence of Cat S confirmed by PCR of the supernatant.

Sf9 cells were infected with recombinant baculovirus at a MOI of 5 for 48-72 hrs. Cell pellet was lysed and incubated in buffer at pH 4.5 at 37 for 2 hours to activate Cat S from pro-form to active mature form (Bromme, D & McGrath, M., Protein Science, 1996, 5:789-791.) Presence of Cat S was confirmed by SDS-PAGE and Western blot using rabbit anti-human proCat S.

Inhibition of Cathepsin S

Human recombinant cathepsin S expressed in Baculovirus is used at a final concentration of 10nM in buffer. Buffer is 50mM Na Acetate, pH 6.5, 2.5mMEDTA, 2.5mMTCEP. Enzyme is incubated with either compound or DMSO for 10 min at 37C. Substrate 7-amino-4-methylcoumarin, CBZ-L-valyl-L-valyl-L-arginineamide (custom synthesis by Molecular Probes) is diluted to 20uM in water (final concentration of 5uM), added to assay and incubated for additional 10 minutes at 37 C. Compound activity is measured by diminished fluorescence compared to DMSO control when read at 360nm excitation and 460nm emission.

Examples listed above were evaluated for inhibition of cathepsin S in the above assay. All had IC₅₀ values of 100 micromolar or below.

Inhibition of Cathepsin K, F, L and B:

Inhibition of these enzymes by particular compounds of the invention may be determined
5 without undue experimentation by using art recognized methods as provided hereinbelow
each of which is incorporated herein by reference:

Cathepsin B, and L assays are to be found in the following references:

- 10 1. Methods in Enzymology, Vol.244, Proteolytic Enzymes: Serine and Cysteine
Peptidases, Alan J. Barrett, ed.

Cathepsin K assay is to be found in the following reference:

- 15 2. Bromme, D., Okamoto, K., Wang, B. B., and Biroc, S. (1996) *J. Biol. Chem.* **271**,
2126-2132.


20 Cathepsin F assays are to be found in the following references:

3. Wang, B., Shi, G.P., Yao, P.M., Li, Z., Chapman, H.A., and Bromme, D. (1998) *J.*
Biol. Chem. **273**, 32000-32008.
- 25 4. Santamaria, I., Velasco, G., Pendas, A.M., Paz, A., and Lopez-Otin, C (1999) *J. Biol.*
Chem. **274**, 13800-13809.

Preferred compounds to be evaluated for inhibition of Cathepsin K, F, L and B in the
30 above assays desirably have IC₅₀ values of 100 micromolar or below.

What is claimed is

1. A compound of formula (I):



wherein:

10 A is -C(O)- or -CH(OR8)-;

15 R1 is alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl or amino wherein R1 is optionally substituted by one or more R_a;

20 R_a is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, alkanoyl, aroyl, alkoxycarbonyl, aryloxycarbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_a may be further optionally substituted by one or more R_b;

30 R_b is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R2 is hydrogen, OR_i or lower alkyl;

R3 is hydrogen or lower alkyl;

40 R4 is hydrogen or lower alkyl;

R5 is hydrogen, alkyl, cycloalkyl, aryl, heterocyclyl or heteroaryl, wherein R5 is optionally substituted by one or more R_c;

5 R_c is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxycarbonyl, aryloxy carbonyl, alkanoyl, aroyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, 10 alkoxycarbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally 15 substituted by one or more R_d;

R_d is selected from the group consisting of alkyl, cycloalkyl, aryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

20 R6 is hydrogen or lower alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

25 R7 is hydrogen, alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, cycloalkyl, aryl, heterocyclyl, aryl, heteroaryl or cyano, wherein R7 is optionally substituted by one or more R_e;

30 R_e is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, alkoxycarbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, heteroarylalkoxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio, arylthio, arylalkylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, 35 alkoxycarbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_e may be further optionally 40 substituted by one or more R_f;

R_f is selected from the group consisting of alkyl, cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, 45 methyl or methoxy, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxycarbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, carbamoyl

wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

or R₆ and R₇ together with the carbon they are attached form a 4 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, imidazolyl and pyridinyl, C1-5 alkanoyl, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or disubstituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_g may be further optionally substituted by one or more R_h;

R_h is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, lower alkyl or lower alkoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranyl and thiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by

alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl and morpholinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl and pyridinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R₈ is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl or arylalkyl;

R_i is hydrogen or lower alkyl;

X is O or S;

and

pharmaceutically acceptable salts, esters or tautomers thereof.

2. The compound according to claim 1

wherein:

R₁ is C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, pyranal and thiopyranal, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl and amino wherein R₁ is optionally substituted by one or more R_a;

R_a is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or

di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl or phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoxyloxy, arylcarbamoxyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, pyrazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl, aryl, C1-8 alkoxy, aryloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R₂ is hydrogen, OR_i or C1-5 alkyl;

R₃ is hydrogen or C1-5 alkyl;

R₄ is hydrogen or C1-5 alkyl;

R₅ is hydrogen, C1-8 alkyl, C3-7 cycloalkyl or aryl wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl,
 heterocyclcyl selected from the group consisting of pyrrolidinyl, piperidinyl,
 morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from
 5 the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl,
 pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl,
 pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl,
 benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl,
 10 quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8
 alkoxy, aryloxy, alkanoyl, aroyl, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8
 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be
 independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclcyl selected
 from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl,
 15 thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group
 consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl,
 isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl,
 pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl,
 benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl,
 20 quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8
 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, ureido wherein either
 nitrogen atom may be independently substituted by alkyl, aryl, heterocyclcyl
 selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl,
 thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group
 consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl,
 25 isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl,
 pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl,
 benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl,
 quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl,
 alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy,
 30 arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl,
 arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono
 or di-substituted by alkyl, aryl, heterocyclcyl selected from the group consisting of
 pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl
 or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl,
 35 oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl,
 triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl,
 indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl,
 purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl,
 phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro,
 40 amidino and guanidino, R_c may be further optionally substituted by one or more
 R_d;

R_d is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl,
 aryl, arylalkyl, C1-8 alkoxy, aryloxy, arylalkoxy, aroyl, amino, halogen,
 45 hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R7 is hydrogen, C1-8 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-7 cycloalkyl, aryl or cyano, wherein R7 is optionally substituted by one or more R_e;

R_e is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, heteroarylC1-8alkoxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, arylC1-8 alkylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_e may be further optionally substituted by one or more R_f;

5 R_f is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, 10 isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, 20 isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, 30 alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, 40 hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

45

or R6 and R7 together with the carbon they are attached form a 4 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g ;

5 R_g is selected from the group consisting of alkyl, cycloalkyl, phenyl, heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, C1-5alkanoyl, aroyl, C1-5alkoxycarbonyl, aryloxy carbonyl, carbamoyl wherein the nitrogen atom may be
10 independantly mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, C1-5 alkylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, C1-5 alkylaminosulfonyl, arylaminosulfonyl, halogen, hydroxy,
15 oxo, carboxy and cyano, R_g may be further optionally substituted by one or more R_h ;

R_h is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, C1-5 alkyl or C1-5 alkoxy, heterocyclyl selected from the group consisting of piperidinyl,
20 morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from
25 the group consisting of piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo, carboxy and cyano; and

R_8 is hydrogen, alkyl, cycloalkyl-alkyl or arylalkyl;
30

R_i is hydrogen or C1-8 alkyl;

X is O or S.
35

3. The compound according to claim 2

40 wherein:

R_1 is C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl,
45 indolinyl, pyranyl and thiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl,

pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl or amino; wherein R₁ is optionally substituted by one or more R_a;

5 R_a is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl
 10 selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5
 15 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl
 20 selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be
 25 independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5
 30 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of
 35 furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_a may be further optionally substituted by one or more R_b;

40 R_b is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R₂ is hydrogen, OR_i or C1-3 alkyl;

45 R₃ is hydrogen or C1-3 alkyl;

R4 is hydrogen or C1-3 alkyl;

R5 is hydrogen, C1-5 alkyl, C3-7 cycloalkyl or aryl wherein R5 is optionally substituted by one or more R_c;

5 R_c is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, C1-5 alkanoyl, aroyl, C1-5 alkoxy carbonyl, aryloxy carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy carbonylamino, aryloxy carbonylamino, C1-5 alkyl carbamoyloxy, aryl carbamoyloxy, C1-5 alkyl sulfonylamino, aryl sulfonylamino, C1-5 alkyl aminosulfonyl, aryl aminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, amidino and guanidino, R_c may be further optionally substituted by one or more R_d;

40 R_d is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, arylalkyl, C1-5 alkoxy, aryloxy, aryl C1-5 alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

45 R6 is hydrogen or C1-8 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S;

5 R₇ is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-7 cycloalkyl, aryl or cyano, wherein R₇ is optionally substituted by one or more R_c;

10 R_c is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxaliny, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, heteroarylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxaliny, C1-5 alkanoylamino, aroylamino, C1-5 alkylthioarylthioarylC1-5 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxaliny, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxaliny, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_f;

45 R_f is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl,

pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

or R6 and R7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl or a heterocyclic ring selected from the group consisting of pyrrolidinyl, piperidinyl, hexahydropyridazinyl, hexahydropyrimidinyl, piperazinyl, pyranyl, thiopyranyl, 1,2-thiazinanyl-1,1-dioxide, 1,2,6-thiadiazinanyl-1,1-dioxide, isothiazolidinyl-1,1-dioxide and imidazolidinyl-2,4-dione, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of alkyl, cycloalkyl, phenyl, heteroaryl selected from the group consisting of furanyl and thienyl, C1-3 alkanoyl, benzoyl, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting

of furanyl, thienyl and pyridinyl, C1-3 alkylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, C1-5 alkylaminosulfonyl, phenylaminosulfonyl, halogen, hydroxy, carboxy and cyano, R_g may be further optionally substituted by one or more R_h ;

5

R_h is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen, C1-3 alkyl and C1-3 alkoxy, piperidinyl, morpholinyl, piperazinyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, benzimidazolyl, quinolinyl, isoquinolinyl, C1-3 alkoxy, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo and cyano.

10

15 R_8 is hydrogen, C1-8 alkyl, C3-6 cycloalkyl-alkyl or aryl-C1-3 alkyl;

R_i is hydrogen or C1-5 alkyl;

20

4. The compound according to claim 3

wherein:

25

R_1 is C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyranyl and thiopyranyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or amino wherein R_1 is optionally substituted by one or more R_a ;

30

35 R_a is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl,

40

45

benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl or aryl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy, cyano and nitro, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, C1-5 alkoxy, aryloxy, halogen, hydroxy, oxo, carboxy and cyano;

R2 is hydrogen, ORi or methyl;

R3 is hydrogen or methyl;

R4 is hydrogen or methyl;

R⁵ is C1-5 alkyl, C3-7 cycloalkyl or phenyl wherein R⁵ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino,

arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, aryl, arylC1-5alkyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy and cyano;

R_6 is hydrogen or C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

R_7 is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-7 cycloalkyl, phenyl, naphthyl or cyano, wherein R_7 is optionally substituted by one or more R_e ;

R_e is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, heteroarylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, arylC1-5 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be

independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen, methyl and methoxy, naphthyl optionally substituted by one or more groups selected from halogen, methyl and methoxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, C1-5 alkoxy, aryloxy, arylC1-5alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano;

or R₆ and R₇ together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl or a

heterocyclic ring selected from the group consisting of piperidinyl, hexahydropyridazinyl, hexahydropyrimidinyl, piperazinyl, pyranyl and thiopyranyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g ;

- 5 R_g is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, C1-3 alkanoyl, benzoyl, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, carboxy and cyano, R_g may be further optionally substituted by one or more R_h ;

- 10 R_h is selected from the group consisting of methyl, phenyl optionally substituted by one or more groups selected from halogen, methyl and methoxy, piperidinyl, morpholinyl, piperazinyl, pyridinyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo and cyano;

- 20 R_8 is hydrogen, C1-5 alkyl, C5-6 cycloalkyl-C1-3 alkyl or benzyl;

R_i is hydrogen or methyl; and

- 25 X is O.

- 30 5. A compound according to claim 4
wherein:

- 35 R_1 is C1-5 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl, piperazinyl, pyranyl and thiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or amino, wherein R_1 is optionally substituted by one or more R_a ;

- 40 R_a is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-3 alkoxy, phenoxy, C1-3 alkoxycarbonyl,

aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, phenyl or naphthyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, aryl, C1-3 alkoxy, phenoxy, halogen, hydroxy, oxo, carboxy and cyano;

R_2 is hydrogen or Ori;

R_3 is hydrogen;

R_4 is hydrogen;

R_5 is C1-5 alkyl, C3-6 cycloalkyl or phenyl, wherein R_5 is optionally substituted by one or more R_c ;

R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, phenoxy, aroyl, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio, phenylthio, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino,

arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of C1-5 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, arylC1-3alkyl, C1-5 alkoxy, phenoxy, arylC1-3alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy and cyano;

R₆ is hydrogen or C1-3alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

R₇ is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, C3-6 cycloalkyl, phenyl or cyano, wherein R₇ is optionally substituted by one or more R_e;

R_e is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, heteroarylC1-3alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio, arylC1-3alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-5 alkoxycarbonylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, and indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen and methyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl and indolyl, C1-5 alkoxy,

aryloxy, arylC1-3alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5
alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be
independently mono or di-substituted by C1-5 alkyl or aryl, C1-5
alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either
5 nitrogen atom may be independently substituted by C1-5 alkyl or aryl, C1-
5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy,
arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5
alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom
may be independently mono or di-substituted by C1-5 alkyl or aryl,
10 halogen, hydroxy, oxo, carboxy and cyano;

or

15 R₆ and R₇ together with the carbon they are attached form a carbocyclic ring selected
from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl or
heterocyclic ring selected from the group consisting of piperidinyl, morpholinyl and
thiomorpholinyl, the carbocyclic or heterocyclic ring being optionally substituted with
one or more R_g;

20 R_g is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, C1-3
alkanoyl, C1-3 alkoxycarbonyl and carbamoyl wherein the nitrogen atom may be
independently mono or disubstituted by C1-5 alkyl; R_g may be further optionally
substituted by one or more R_h; and

25 R_h is phenyl or amino wherein the nitrogen atom may be independently mono or
di-substituted by C1-3 alkyl and

R₈ is hydrogen or methyl.

30 6. The compound according to claim 5

wherein:

35 R₁ is C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group
consisting of piperidinyl, morpholinyl, piperazinyl, pyranyl and thiopyranyl; heteroaryl
selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl,
40 imidazolyl, pyridinyl and indolyl or amino, wherein R₁ is optionally substituted by one
or more R_a;

45 R_a is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl,
heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and
piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl,
pyrrolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl,

benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-3 alkoxy, C1-3 alkoxy carbonyl, aryloxy carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl or heteroaryl selected from the group consisting of pyrrolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl and benzthiazolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or phenyl, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of pyrrolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, C1-3 alkoxy, halogen and hydroxy;

R_2 is hydrogen;

R_5 is C1-5 alkyl, C5-6 cycloalkyl or phenyl wherein R_5 is optionally substituted by one or more R_c ;

R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, 4-piperidinyl, 4-morpholinyl, piperazinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-5 alkoxy, phenoxy, aroyl, C1-5 alkoxy carbonyl, aryloxy carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, C1-5 alkanoylamino, C1-3 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or phenyl, C1-5 alkoxy carbonylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of C1-5 alkyl, C5-6 cycloalkyl, phenyl, benzyl, C1-5 alkoxy, phenoxy, benzyloxy, aroyl, halogen, hydroxy, oxo, carboxy and cyano;

R_6 is hydrogen or C1-3 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N, O and S;

R7 is hydrogen, C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S, phenyl or cyano, wherein R7 is optionally substituted by one or more R_e;

R_e is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, benzyloxy, pyridylC1-3alkoxy, thienylC1-3alkoxy, furanylC1-3alkoxy, C1-3 alkoxycarbonyl, phenoxyoxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, methylthio, benzylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-3 alkoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of C1-3 alkyl, phenyl optionally substituted by one or more groups selected from the group consisting of halogen and methyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl and pyridinyl, C1-3 alkoxy, aryloxy, benzyloxy, C1-5 alkoxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-3 alkylsulfonylamino, arylsulfonylamino, C1-3 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano;

or R6 and R7 together with the carbon they are attached form cyclopropyl or a heterocyclic ring selected from the group consisting of piperidinyl, pyranyl and thiopyranyl, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g; and

R_g is selected from the group consisting of methyl, benzyl, acetyl, benzoyl, benzyloxycarbonyl and ethoxycarbonyl; and

R8 is hydrogen.

7. The compound according to claim 6

wherein:

5 R1 is C3-6 cycloalkyl, phenyl, naphthyl, piperidinyl, morpholinyl, piperazinyl, pyranyl, thiopyranyl, furanyl, thienyl, pyrrolyl or amino, wherein R1 is optionally substituted by one or more R_a;

10 R_a is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5
15 alkanoylamino, aroylamino, ureido wherein either nitrogen atom may be independently substituted by C1-3alkyl or phenyl, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-3alkyl, C1-3alkoxy, halogen and hydroxy;

20 R5 is C1-5 alkyl, C5-6 cycloalkyl or phenyl, wherein R5 is optionally substituted by one or more groups of the formula R_c;

25 R_c is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, 4-piperidinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, C1-5 alkoxycarbonyl, C1-5 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, C1-3 alkylthio, C1-3 alkoxycarbonylamino, C1-3 alkylsulfonylamino, amino wherein the nitrogen atom may be independently
30 mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

35 R_d is selected from the group consisting of C1-3 alkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, benzyloxy, benzoyl, halogen, hydroxy, oxo, carboxy and cyano;

wherein the configuration at the stereocenter defined by R4 and R5 and the carbon they are attached to is defined as L;

40 R6 is hydrogen or C1-2 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S;

45 R7 is C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S; phenyl or cyano wherein R7 is optionally substituted by one or more groups of the formula R_e;

R_e is selected from the group consisting of methyl, C3-6 cycloalkyl, phenyl, naphthyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, benzyloxy, pyridylC1-3alkoxy, thienylC1-3alkoxy, furanylC1-3alkoxy, C1-5 alkanoylamino, aroylamino, methylthio, benzylthio, C1-3 alkoxy-carbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f .

R_f is selected from the group consisting of C1-3 alkyl, phenyl optionally substituted by one or more groups selected from halogen and methyl, C1-3 alkoxy, aryloxy, benzyloxy, C1-3 alkoxy-carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano;

or R_6 and R_7 together with the carbon they are attached form cyclopropyl or a heterocyclic ring selected from the group consisting of piperidinyl and pyranyl, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g ; and

R_g is methyl.

8. The compound according to claim 7

wherein:

R_1 is cyclopropyl, cyclohexyl, phenyl, naphthyl, piperidinyl, morpholinyl, piperazinyl, pyranyl, thiopyranyl or amino wherein R_1 is optionally substituted by one or more R_a ;

R_a is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxy-carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-3 alkoxy, halogen and hydroxy,

R5 is C1-5 alkyl or C5-6 cycloalkyl wherein R5 is optionally substituted by one or more R_c;

5 R_c is selected from the group consisting of methyl, C3-6 cycloalkyl, phenyl, naphthyl, thienyl, imidazolyl, pyridinyl, indolyl, C1-4 alkoxy, C1-5 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

10 R_d is selected from the group consisting of methyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, halogen and hydroxy;

R7 is C1-5 alkyl or phenyl, wherein R7 is optionally substituted by one or more R_e;

15 R_e is selected from the group consisting of C3-6 cycloalkyl, phenyl, naphthyl, thienyl, imidazolyl, pyridinyl, indolyl, methoxy, benzyloxy, C1-3 alkanoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, benzylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, methoxycarbonylamino, amino wherein the nitrogen atom may be independently
20 mono or di-substituted by C1-3 alkyl, halogen, hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of methyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, methoxy, phenoxy, benzyloxy, methoxycarbonyl, amino wherein the
25 nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy and carboxy;

or R6 and R7 together with the carbon they are attached form a cyclopropyl ring.

30

9. The compound according to claim 8

35 wherein:

40 R1 is cyclopropyl, cyclohexyl, phenyl, naphthyl, piperidiny, morpholinyl, piperazinyl, pyranly or thiopyranly, wherein R1 is optionally substituted by one or more R_a;

R_a is selected from the group consisting of pyrrolyl, imidazolyl, indolyl, benzimidazolyl, methoxy, methoxycarbonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy
45 and carboxy, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of methoxy, halogen and hydroxy;

R_c is selected from the group consisting of methyl, C3-6 cycloalkyl, phenyl, naphthyl, C1-4 alkoxy, C1-3 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of methyl, phenyl, methoxy, halogen and hydroxy;

R_6 is hydrogen;

R_e is selected from the group consisting of C5-6 cycloalkyl, phenyl, naphthyl, thienyl, indolyl, methoxy, benzyloxy, methylthio, benzylthio, methoxycarbonylamino, halogen, hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f ; and

R_f is selected from the group consisting of methyl, phenyl optionally substituted by halogen, methoxy, phenoxy, benzyloxy, methoxycarbonyl, halogen, hydroxy and carboxy.

10. The compound according to claim 9

wherein:

R_1 is 4-morpholinyl, 4-pyranyl or phenyl;

R_5 is C1-5 alkyl wherein R_5 is optionally substituted by one or more R_c ;


R_c is selected from the group consisting of C3-6 cycloalkyl, phenyl and 2-naphthyl, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of methyl and halogen;

R_e is selected from the group consisting of C5-6 cycloalkyl, phenyl, naphthyl, indolyl, benzyloxy, methylthio, benzylthio, halogen and carboxy, R_e may be further optionally substituted by one or more R_f ; and

R_f is selected from the group consisting of methyl, methoxy, methoxycarbonyl, halogen and hydroxy.

11. A compound of formula (Ia):



wherein:

A is -C(O)- or -CH(OR8)-;

R1 is alkyl, alkoxy, cycloalkyl, aryl, aryloxy, benzyloxy, heterocyclyl, heteroaryl or amino wherein R1 is optionally substituted by one or more Ra;

Ra is selected from the group consisting of a bond, alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, alkanoyl, aroyl, alkoxy carbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxy carbonylamino, aryloxy carbonylamino, alkyl carbamoyloxy, aryl carbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; Ra may be further optionally substituted by one or more Rb;

Rb is selected from the group consisting of alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, acyl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R2 is hydrogen, ORj or lower alkyl;

R3 is hydrogen or lower alkyl;

R4 is hydrogen or lower alkyl;

R5 is hydrogen, alkyl, cycloalkyl, aryl, heterocyclyl or heteroaryl, wherein R5 is optionally substituted by one or more R_c;

5 R_c is selected from the group consisting of a bond, alkyl, cycloalkyl, bicycloalkyl, aryl, indenyl, indanyl, dihydronaphthyl, tetrahydronaphthyl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxycarbonyl, aryloxy carbonyl, alkanoyl, aroyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxycarbonylamino, aryloxy carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_d;

20 R_d is selected from the group consisting of alkyl, cycloalkyl, aryl, arylalkyl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, amino, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

25 or R4 and R5 together with the carbon they are attached form a 3 to 7 membered carbocyclic ring optionally substituted with one or more R_d;

30 R6 is hydrogen or lower alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S;

35 R7 is hydrogen, alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S, cycloalkyl, aryl, heterocyclyl, heteroaryl or cyano, wherein R7 is optionally substituted by one or more R_c;

40 R_e is selected from the group consisting of a bond alkyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkanoyl, aroyl, alkoxycarbonyl, aryloxy carbonyl, alkanoyloxy, aroyloxy, heteroarylalkoxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylalkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be

45

independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxy-carbonylamino, aryloxy-carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_f,

R_f is selected from the group consisting of alkyl, cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, methyl or methoxy, heterocyclyl, heteroaryl, alkoxy, aryloxy, arylalkoxy, alkoxy-carbonyl, aryloxy-carbonyl, alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkanoylamino, aroylamino, alkylcarbamoyl, arylcarbamoyl, alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl or heteroaryl, alkoxy-carbonylamino, aryloxy-carbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl or heteroaryl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

or R₆ and R₇ together with the carbon they are attached form a 3 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of a bond, alkyl, cycloalkyl, bicycloalkyl, benzyl, aryl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, tetrahydrothiopyranyl, indolinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrimidinyl, pyrrolyl, imidazolyl and pyridinyl, C1-5 alkanoyl, aroyl, C1-5 alkoxy-carbonyl, aryloxy-carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or disubstituted by alkyl, cycloalkyl, bicycloalkyl, aryl, aroyl, benzoyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or

heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_g may be further optionally substituted by one or more R_h ;

R_h is selected from the group consisting of a bond, C1-5 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, lower alkyl or lower alkoxy, benzyl, benzyloxy, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl and morpholinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl and pyridinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino; R_h may be further optionally substituted by one or more R_i ;

R_i is alkyl, hydroxy, oxo and halogen;

R_j is hydrogen, alkyl, cycloalkyl;

R_8 is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl or arylalkyl;

X is O or S;

and

pharmaceutically acceptable salts, esters or tautomers thereof.

12. The compound according to claim 11

wherein:

R1 is C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, 2-oxa-5-aza-bicyclo[2,2,1]heptanyl, piperazinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl and amino wherein R1 is optionally substituted by one or more R_a;

R_a is selected from the group consisting of a bond, C1-8 alkyl, C3-7 cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, pyrazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl,

pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinoliny, isoquinoliny, quinazoliny, quinoxaliny, carbazolyl, phenothiaziny and phenoxaziny, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino, guanidino; R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl, aryl, heteroaryl, C1-8 alkoxy, aryloxy, alkanoyl, aroyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, cycloalkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indoliny, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinoliny and isoquinoliny, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

R_2 is hydrogen, OR₃ or C1-3 alkyl;

R_3 is hydrogen or C1-5 alkyl;

R_4 is hydrogen or C1-5 alkyl;

R_5 is hydrogen, C1-8 alkyl, C3-7 cycloalkyl, aryl wherein R_5 is optionally substituted by one or more R_c ;

R_c is selected from the group consisting of a bond, C1-8 alkyl, C3-7 cycloalkyl, bicycloalkyl, aryl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indoliny; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinoliny, isoquinoliny, quinazoliny, quinoxaliny, carbazolyl, phenothiaziny and phenoxaziny, C1-8 alkoxy, aryloxy, alkanoyl, aroyl, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indoliny, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl,

5 benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl,
 quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8
 alkanoylamino, aroylamino, C1-8 alkylthio wherein the sulfur atom may be
 oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be
 10 oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be
 independently substituted by alkyl, aryl, heterocyclyl selected from the group
 consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl
 and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl,
 pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl,
 15 oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl,
 pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl,
 benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl,
 carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino,
 C1-5aryloxycarbonylamino, alkylcarbamoxyloxy, arylcarbamoxyloxy,
 20 alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl,
 amino wherein the nitrogen atom may be independently mono or di-substituted by
 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl,
 piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl
 selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl,
 25 thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl,
 tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl,
 isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl,
 quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl
 and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and
 guanidino, R_c may be further optionally substituted by one or more R_d ;

30 R_d is selected from the group consisting of C1-8 alkyl, C3-6 cycloalkyl,
 aryl, arylalkyl, C1-8 alkoxy, aryloxy, arylalkoxy, aroyl, amino, halogen,
 hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino;

35 R_7 is hydrogen, C1-8 alkyl wherein one or more carbon atoms are optionally replaced by
 1 to two heteroatoms selected from the group consisting of N, O and S, C3-7 cycloalkyl,
 aryl, heteroaryl or cyano, wherein R_7 is optionally substituted by one or more R_c ;

40 R_c is selected from the group consisting of a bond, C1-8 alkyl, C3-7 cycloalkyl,
 aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl,
 morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from
 the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl,
 pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl,
 45 pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl,
 benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl,
 quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8
 alkoxy, aryloxy, arylC1-8alkoxy, heteroarylC1-8alkoxy wherein the heteroaryl is
 as hereinabove described in this paragraph, C1-8 alkoxycarbonyl, aryloxycarbonyl,
 C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be

independently mono or di-substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylC1-8 alkylthio, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino and guanidino, R_c may be further optionally substituted by one or more R_f,

R_f is selected from the group consisting C1-8 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from the group consisting halogen, methyl and methoxy, heterocyclyl selected from the group consisting pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkoxy, aryloxy, arylC1-8alkoxy, C1-8 alkoxycarbonyl, aryloxycarbonyl, C1-8 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-

substituted by C1-8 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, C1-8 alkanoylamino, aroylamino, C1-8 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, alkoxycarbonylamino, aryloxycarbonylamino, alkylcarbamoyloxy, arylcarbamoyloxy, alkylsulfonylamino, arylsulfonylamino, alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl and indolinyl, or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, carbazolyl, phenothiazinyl and phenoxazinyl, halogen, hydroxy, oxo, carboxy, cyano, nitro, amidino, and guanidino;

or R6 and R7 together with the carbon they are attached form a 3 to 7 membered carbocyclic or heterocyclic ring, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

R_g is selected from the group consisting of a bond, alkyl, cycloalkyl, bicycloalkyl, phenyl, naphthyl, benzyl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, tetrahydrothiopyranyl, indolinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrimidinyl and pyridinyl, C1-5alkanoyl, aroyl, C1-5alkoxycarbonyl, aryloxycarbonyl, carbamoyl

wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl, aryl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, thiazolyl, imidazolyl, pyridinyl, benzimidazolyl and quinolinyl, amino
5 wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, benzyl, benzoyl or naphthoyl, C1-5 alkylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidised to a sulfoxide or sulfone, C1-5 alkylaminosulfonyl, arylaminosulfonyl, halogen, hydroxy, oxo, carboxy and cyano, R_g may be further
10 optionally substituted by one or more R_h ;

R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, aryl optionally substituted by one or more groups selected from halogen, C1-5 alkyl or C1-5 alkoxy, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, indolyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group
15 consisting of furanyl, thienyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, benzimidazolyl, quinolinyl, isoquinolinyl, quinazolinyl, benzoxazolyl and quinoxalinyl, C1-5 alkoxy, aryloxy, amino wherein the nitrogen atom may be independently mono or di-substituted by alkyl, aryl, heterocyclyl selected from the group consisting of piperidinyl and morpholinyl, or heteroaryl selected from the group consisting of furanyl, thienyl and pyridinyl, halogen, hydroxy, oxo, carboxy and cyano; and
20

25 R_8 is hydrogen, alkyl, cycloalkyl-alkyl or arylalkyl;

R_i is C1-8 alkyl, hydroxy, oxo and halogen

30 R_j is hydrogen or alkyl and

X is O.

35 13. A compound according to claim 12

wherein:

40 A is $-C(O)-$;

R_1 is C1-5 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, 2-oxa-5-aza-bicyclo[2,2,1]heptanyl, piperazinyl, tetrahydropyranyl and tetrahydrothiopyranyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl,
45

thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or amino, wherein R₁ is optionally substituted by one or more R_a;

5 R_a is selected from the group consisting of a bond, C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, tetrazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, 10 quinolinyl and isoquinolinyl, C1-3 alkoxy, phenoxy, C1-3 alkoxycarbonyl, aryloxy, carbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 15 alkanoylamino, aroylamino, C1-5 alkylthio, arylthio, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl, phenyl or naphthyl; C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be 20 independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, 25 halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

30 R_b is selected from the group consisting of C1-3 alkyl, C5-6 cycloalkyl, aryl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-3 alkoxy, phenoxy, C1-8 alkanoyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 35 cycloalkyl, phenyl or naphthyl, halogen, hydroxy, oxo, carboxy and cyano;

R₂ is hydrogen;

40 R₃ is hydrogen;

R₄ is hydrogen;

45 R₅ is C1-5 alkyl, C3-6 cycloalkyl or phenyl, wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of a bond, C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl, heteroaryl selected from the group consisting of furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, phenoxy, aroyl, C1-5 alkoxycarbonyl, aryloxy, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, phenylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl, phenyl or heteroaryl selected the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkoxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, phenyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d ;

R_d is selected from the group consisting of C1-5 alkyl, C5-6 cycloalkyl, phenyl, naphthyl, arylC1-3alkyl, C1-5 alkoxy, phenoxy, arylC1-3alkoxy, aroyl, amino, halogen, hydroxy, oxo, carboxy and cyano;

R_6 is hydrogen or C1-3alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S;

R_7 is hydrogen, C1-5 alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N, O and S, C3-6 cycloalkyl, phenyl, naphthyl, heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl or cyano, wherein R_7 is optionally substituted by one or more R_e ;

R_e is selected from the group consisting of C1-3 alkyl, C3-6 cycloalkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, quinolinyl and isoquinolinyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, heteroarylC1-3alkoxy

wherein the heteroaryl is as hereinabove described in this paragraph, C1-5 alkoxy carbonyl, aryloxy carbonyl, C1-5 alkanoyloxy, aryloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl and indolyl, C1-5 alkanoylamino, aroylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylC1-3alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or phenyl, C1-5 alkoxy carbonylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, phenyl, naphthyl, heterocyclyl selected from the group consisting of piperidinyl, morpholinyl and piperazinyl or heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl and pyridinyl, indolyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_g.

R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, morpholinyl and piperazinyl; heteroaryl selected from the group consisting of furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl and indolyl, C1-5 alkoxy, aryloxy, arylC1-3alkoxy, C1-5 alkoxycarbonyl, aryloxycarbonyl, C1-5 alkanoyloxy, aroyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, C1-5 alkanoylamino, aroylamino, C1-5 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, arylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, ureido wherein either nitrogen atom may be independently substituted by C1-5 alkyl or aryl, C1-5 alkoxycarbonylamino, aryloxycarbonylamino, C1-5 alkylcarbamoyloxy, arylcarbamoyloxy, C1-5 alkylsulfonylamino, arylsulfonylamino, C1-5 alkylaminosulfonyl, arylaminosulfonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or aryl, halogen, hydroxy, oxo, carboxy and cyano;

or

R6 and R7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or heterocyclic ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, tetrahydropyrany and tetrahydrothiopyrany, the carbocyclic or heterocyclic ring being optionally substituted with one or more R₄;

R_g is selected from the group consisting of a bond, C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, heterocyclyl selected from the group consisting of pyrrolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl and morpholinyl, heteroaryl selected from the group consisting of thienyl, pyrimidinyl and pyridinyl, C1-5 alkanoyl, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or disubstituted by C1-5 alkyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, benzyl, benzoyl or naphthoyl, R_g may be further optionally substituted by one or more R_h ; and

R_h is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, phenyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of piperidinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of thienyl, pyrrolyl, pyridinyl, pyrimidinyl, indolyl, benzimidazolyl, quinolinyl and isoquinolinyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, oxo, cyano and trifluoromethyl.

14. The compound according to claim 13

wherein:

R_1 is C3-6 cycloalkyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, tetrahydropyranyl, furanyl, thienyl, pyrrolyl or amino, wherein R_1 is optionally substituted by one or more R_a ;

R_a is selected from the group consisting of a bond, C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl; C1-5 alkoxy carbonylamino, C1-5 alkanoylamino, aroylamino, ureido wherein either nitrogen atom may be independently substituted by C1-3 alkyl or phenyl; C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b ;

R_b is selected from the group consisting of C1-3 alkyl, C1-3 alkoxy, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, oxo and hydroxy;

R5 is C1-5 alkyl, C3-6 cycloalkyl or phenyl, wherein R5 is optionally substituted by one or more groups of the formula R_c;

5 R_c is selected from the group consisting of a bond, C1-3 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, indenyl, indanyl, 4-piperidinyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, C1-5 alkoxycarbonyl, C1-5 alkanoyloxy, benzoyloxy, carbamoyl wherein the nitrogen atom may be independently mono or di-
10 substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, C1-3 alkylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-3 alkoxycarbonylamino, C1-3 alkylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

15 R_d is selected from the group consisting of C1-3 alkyl, phenyl, benzyl, C1-3 alkoxy, phenoxy, benzyloxy, benzoyl, halogen, hydroxy, oxo, carboxy and cyano;

20 wherein the configuration at the stereocenter defined by R4 and R5 and the carbon they are attached to is defined as L;

R6 is hydrogen or C1-2alkyl one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S;

25 R7 is C1-5 alkyl said alkyl being optionally interrupted by 1 to two heteroatoms selected from the group consisting of N,O and S; phenyl, pyridinyl or cyano wherein R7 is optionally substituted by one or more groups of the formula R_e;

30 R_e is selected from the group consisting of a bond, methyl, C3-6 cycloalkyl, phenyl, naphthyl, furanyl, thienyl, thiazolyl, imidazolyl, pyridinyl, pyrrolidinyl, piperidinyl, indolyl, C1-3 alkoxy, benzyloxy, pyridinylC1-3alkoxy, thienylC1-3alkoxy, furanylC1-3alkoxy, C1-5 alkanoylamino, aroylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, benzylthio
35 wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, C1-3 alkoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

40 R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, C1-3 alkoxy, aryloxy, benzyloxy, C1-3 alkoxycarbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-
45 substituted by C1-3 alkyl or phenyl, C1-5 alkanoylamino, aroylamino, amino wherein the nitrogen atom may be independently mono or di-

substituted by C1-3 alkyl or phenyl, halogen, hydroxy, oxo, carboxy and cyano;

5 R6 and R7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclopentyl, cyclohexyl and cycloheptyl or heterocyclic ring selected from the group consisting of pyrrolidinyl, piperidinyl and tetrahydropyranyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

10 R_g is selected from the group consisting of a bond C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, dihydronaphthyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from the group consisting of piperidinyl, tetrahydropyranyl, and tetrahydrothiopyranyl, C1-5 alkanoyl, C1-3 alkoxy carbonyl, carbamoyl
15 wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, benzyl, benzoyl or naphthoyl, R_g may be further optionally substituted by one or more R_h; and

20 R_h is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, phenyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of piperidinyl, indolinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of thienyl, pyridinyl, indolyl, pyrrolyl and benzimidazolyl, amino wherein the nitrogen atom may be independently mono or di-substituted by
25 methyl, halogen, hydroxy, oxo, cyano and trifluoromethyl.

30 15. The compound according to claim 14

wherein:

35 R1 is cyclohexyl, phenyl, naphthyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, or amino wherein R1 is optionally substituted by one or more R_a;

40 R_a is selected from the group consisting of a bond, C1-3 alkyl, C5-6 cycloalkyl, phenyl, furanyl, thienyl, pyrrolyl, imidazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzthiazolyl, C1-3 alkoxy, C1-3 alkoxy carbonyl, carbamoyl wherein the nitrogen atom may be independently mono or di-substituted by C1-3
45 alkyl or phenyl, C1-5 alkoxy carbonylamino, C1-5 alkanoylamino, aroylamino, C1-5 alkylsulfonylamino, arylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, oxo, carboxy and cyano, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of C1-3 alkoxy, halogen and hydroxy,

5

R₅ is C1-5 alkyl wherein R₅ is optionally substituted by one or more R_c;

10

R_c is selected from the group consisting of a bond, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, indanyl, thienyl, imidazolyl, pyridinyl, indolyl, C1-3 alkoxy, C1-5 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

15

R_d is selected from the group consisting of C1-3 alkyl, phenyl, benzyl, methoxy, phenoxy, benzyloxy, benzoyl, halogen and hydroxy;

R₇ is C1-5 alkyl or phenyl, wherein R₇ is optionally substituted by one or more R_e;

20

R_e is selected from the group consisting of a bond, C3-6 cycloalkyl, phenyl, naphthyl, thienyl, imidazolyl, pyrrolidinyl, piperidinyl, pyridinyl, indolyl, C1-5 alkoxy, benzyloxy, C1-3 alkanoylamino, methylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, benzylthio wherein the sulfur atom may be oxidized to a sulfoxide or sulfone, methoxycarbonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, halogen, hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f;

25

30

R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by one or more groups selected from halogen or methyl, methoxy, phenoxy, benzyloxy, methoxycarbonyl, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl or phenyl, halogen, hydroxy and carboxy;

35

R₆ and R₇ together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl and cyclohexyl or heterocyclic ring selected from the group consisting of pyrrolidinyl and piperidinyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

40

R_g is selected from the group consisting of a bond, C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, tetrahydronaphthyl, indanyl, heterocyclyl selected from the group consisting of tetrahydropyranyl and tetrahydrothiopyranyl, acetyl, methoxycarbonyl, carbamoyl wherein the nitrogen atom may be independantly mono or disubstituted by C1-3 alkyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, benzyl or benzoyl, R_g may be further optionally substituted by one or more R_h; and

45

R_h is selected from the group consisting of C1-8 alkyl, C3-7 cycloalkyl, phenyl, benzyl, benzyloxy, heterocyclyl selected from the group consisting of piperidinyl, tetrahydropyranyl and tetrahydrothiopyranyl, heteroaryl selected from the group consisting of thienyl, pyridinyl, indolyl and benzimidazolyl, amino, methylamino, dimethylamino, halogen, hydroxy, oxo, cyano and trifluoromethyl.

16. The compound according to claim 15

wherein:

R₁ is phenyl, naphthyl, pyrrolidinyl, thiomorpholinyl or morpholinyl, wherein R₁ is optionally substituted by one or more R_a;

R_a is selected from the group consisting of a bond, C1-3 alkyl, pyrrolyl, imidazolyl, indolyl, benzimidazolyl, methoxy, methoxycarbonyl, t-butoxycarbonylamino, C1-3 alkylsulfonylamino, amino wherein the nitrogen atom may be independently mono or di-substituted by C1-3 alkyl, halogen, hydroxy, cyano and carboxy, R_a may be further optionally substituted by one or more R_b;

R_b is selected from the group consisting of methoxy, halogen and hydroxy;

R_c is selected from the group consisting of a bond, methyl, C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, tetrahydronaphthyl, indanyl, C1-4 alkoxy, C1-3 alkanoylamino, methylthio, halogen, hydroxy, oxo, carboxy and cyano, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of C1-3 alkyl, phenyl, methoxy, halogen and hydroxy;

R₆ is hydrogen;

R_e is selected from the group consisting of a bond, C5-6 cycloalkyl, phenyl, C1-5 alkoxy, C1-5 alkylamino, pyrrolidinyl, piperidinyl, naphthyl, thienyl, indolyl, methoxy, benzyloxy, methylthio, benzylthio, methoxycarbonylamino, halogen, hydroxy, carboxy and cyano, R_e may be further optionally substituted by one or more R_f; and

R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted by halogen, methoxy, phenoxy, benzyloxy, methoxycarbonyl, halogen, hydroxy and carboxy.

R6 and R7 together with the carbon they are attached form a cyclopropyl ring or heterocyclic ring selected from the group consisting of pyrrolidinyl and piperidinyl, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g;

5 R_g is selected from the group consisting of a bond, C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, α-tetrahydronaphthyl, α-indanyl, tetrahydropyranyl, tetrahydrothiopyranyl and amino wherein the nitrogen atom may be independently mono or di-substituted by C1-5 alkyl, C3-6 cycloalkyl, benzyl or benzoyl, R_g may be further optionally substituted by one or more R_h; and

10 R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, benzyl, benzyloxy, tetrahydropyranyl, heteroaryl selected from the group consisting of thienyl, pyridinyl, indolyl, pyrrolyl and benzimidazolyl, halogen, hydroxy, oxo, cyano or trifluoromethyl, may be further substituted by R_i; and

15 R_i is C1-8 alkyl and halogen.

20 17. The compound according to claim 16

wherein:

25 R1 is phenyl, morpholinyl, thiomorpholinyl or pyrrolidinyl, R1 is optionally substituted by one or more R_a;

R_a is selected from the group consisting of C1-3 alkyl, methoxy, C1-3 alkylsulfonylamino, halogen, hydroxy, cyano and trifluoromethyl;

30 R5 is C1-3 alkyl wherein R5 is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C3-6 cycloalkyl, C7-8 bicycloalkyl, phenyl, naphthyl, β-tetrahydronaphthyl or β-indanyl, R_c may be further optionally substituted by one or more R_d;

35 R_d is selected from the group consisting of methyl, methoxy and halogen;

R7 is C1-5 alkyl wherein R7 is optionally substituted by one or more R_e;

40 R_e is selected from the group consisting of phenyl, naphthyl, C1-5 alkoxy, C1-5 alkylamino, pyrrolidinyl and piperidinyl, R_e may be further optionally substituted by one or more R_f; and

45 R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted with halogen, halogen and hydroxy.

or R6 and R7 together with the carbon they are attached form a cyclopropyl ring or heterocyclic ring selected from the group consisting of piperidinyl and pyrrolidinyl, the cyclopropyl or heterocyclic ring being optionally substituted with one or more R_g; and

R_g is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, α -tetrahydronaphthyl, α -indanyl, tetrahydropyranyl, tetrahydrothiopyranyl and amino group wherein the nitrogen atom may be independantly mono or disubstituted by C1-5 alkyl or C3-6 cycloalkyl; R_g may be further optionally substituted by one or more R_h; and

R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, benzyl, benzyloxy, thienyl, pyrrolyl, indolyl, pyridinyl, halogen, hydroxy, cyano and trifluoromethyl; R_h may be further optionally substituted by one or more R_i; and

R_i is C1-5 alkyl and halogen.

18. The compound according to claim 17

wherein:

R1 is 4-morpholinyl or pyrrolidinyl, wherein R1 is optionally substituted by one or more R_a;

R_a is C1-3 alkylsulfonylamino;

R5 is C1-3 alkyl wherein R5 is optionally substituted by one or more R_c;

R_c is selected from the group consisting of C5-6 cycloalkyl, C7-8 bicycloalkyl, β -tetrahydronaphthyl and β -indanyl, R_c may be further optionally substituted by one or more R_d;

R_d is selected from the group consisting of methyl, methoxy and halogen;

R7 is C1-5 alkyl wherein R7 is optionally substituted by one or more R_e;

R_e is selected from the group consisting of C1-5 alkoxy, C1-5 alkylamino, pyrrolidinyl and piperidinyl, R_e may be further optionally substituted by one or more R_f; and

R_f is selected from the group consisting of C1-5 alkyl, C3-6 cycloalkyl, phenyl optionally substituted with halogen, halogen and hydroxy.

5 or R₆ and R₇ together with the carbon they are attached form heterocyclic ring selected from the group consisting of piperidinyl and pyrrolidinyl, the heterocyclic ring being optionally substituted with one or more R_g; and

10 R_g is selected from the group consisting of C1-8 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl, α -tetrahydronaphthyl, α -indanyl, tetrahydropyranyl and tetrahydrothiopyranyl; R_g may be further optionally substituted by one or more R_h; and

R_h is selected from the group consisting of C1-5 alkyl, C3-7 cycloalkyl, phenyl, pyridinyl, indolyl, thienyl, halogen, hydroxy, cyano and trifluoromethyl; R_h may be further optionally substituted by one or more R_i;

15

R_i is selected from the group consisting of C1-3 alkyl and halogen.

20 19. The compound according to claim 18

20

wherein:

R₁ is 4-morpholinyl;

25 R₅ is C1-3 alkyl wherein R₅ is optionally substituted by one or more R_c;

R_c is selected from the group consisting of cyclohexyl, β -tetrahydronaphthyl and β -indanyl, R_c may be further optionally substituted by one or more R_d;

30

R_d is selected from the group consisting of methyl and halogen;

R₇ is C1-5 alkyl wherein R₇ is optionally substituted by one or more R_e;

35

R_e is selected from the group consisting of C1-2alkoxy and C1-2 alkylamino, R_e may be further optionally substituted by one or more R_f;

R_f is selected from the group consisting of C1-3 alkyl, phenyl, halogen and hydroxy;

or R6 and R7 together with the carbon they are attached form a heterocyclic ring selected from the group consisting of piperidinyl and pyrrolidinyl, the heterocyclic ring being optionally substituted with one or more R_g; and

- 5 R_g is selected from the group consisting of C1-5 alkyl, C3-8 cycloalkyl, C7-8 bicycloalkyl, benzyl and α -tetrahydronaphthyl; R_g may be further optionally substituted by one or more R_h; and

- R_h is C1-5 alkyl, C3-7 cycloalkyl, thienyl, pyridinyl, indolyl, halogen, hydroxy and
10 trifluoromethyl; R_h may be further optionally substituted by one or more R_i;

R_i is methyl.

- 15 20. The compound according to claim 11 wherein:

A is -C(O)-;

- 20 R1 is aryl or aryloxy wherein R1 is optionally substituted by one or more R_a;

R2 is hydrogen;

R3 is hydrogen;

- 25 R4 is hydrogen or lower alkyl;

R5 is alkyl optionally substituted by one or more R_c;

- 30 or R4 and R5 together with the carbon they are attached form a 3 to 7 membered carbocyclic ring optionally substituted with one or more R_d;

R6 is hydrogen or lower alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S;

- 35 R7 is hydrogen, alkyl wherein one or more carbon atoms are optionally replaced by 1 to two heteroatoms selected from the group consisting of N,O and S, cycloalkyl, aryl,

heterocyclyl, heteroaryl or cyano, wherein R7 is optionally substituted by one or more R_e;

5 R6 and R7 together with the carbon they are attached form a carbocyclic ring selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or heterocyclic ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, tetrahydropyranyl and tetrahydrothiopyranyl, the carbocyclic or heterocyclic ring being optionally substituted with one or more R_g;

10

21. The compound according to claim 20 wherein:

R1 is naphthyl, benzyloxy or phenoxy wherein R1 is optionally substituted by one or more R_a;

15

R4 is hydrogen or lower alkyl;

R5 is lower alkyl optionally substituted by one or more R_c;

20 or R4 and R5 together with the carbon they are attached form a 5 to 6 membered carbocyclic ring optionally substituted with one or more R_d;

25 R6 is hydrogen or lower alkyl wherein one or more carbon atoms are optionally replaced by N;

30 R7 is lower alkyl wherein one or more carbon atoms are optionally replaced by heteroatoms selected from the group consisting of N and O, R7 is optionally substituted by one or more R_e;

35 R6 and R7 together with the carbon they are attached form a heterocyclic ring selected from the group consisting of pyrrolidiny and piperidiny each being optionally substituted with one or more R_g.

35

22. The compound according to claim 21 wherein:

40 R4 is hydrogen or lower alkyl;

R5 is C1-4 alkyl optionally substituted by one or more R_c;

or R4 and R5 together with the carbon they are attached form a cyclohexyl ring
optionally substituted with one or more R_d;

R7 is lower alkyl wherein one or more carbon atoms are optionally replaced by
heteroatoms selected from the group consisting of N and O or benzyloxy;

R6 and R7 together with the carbon they are attached form pyrrolidinyl optionally
substituted with one or more R_g.

23. A compound selected from the group consisting of:

N-(Benzyloxymethyl-cyano-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-
oxo-butyramide;

N-[2-(4-Chloro-benzyloxy)-1-cyano-ethyl]-2-cyclohexylmethyl-4-morpholin-4-
yl-4-oxo-butyramide;

N-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-
4-yl-4-oxo-butyramide;

N-(3-Cyano-1-cyclohexylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-
morpholin-4-yl-4-oxo-butyramide;

N-[3-Cyano-1-(2,6-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-
morpholin-4-yl-4-oxo-butyramide;

N-[3-Cyano-1-(3,5-difluoro-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-
morpholin-4-yl-4-oxo-butyramide;

N-[3-Cyano-1-(3-trifluoromethyl-benzyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-cyano-1-(3,3-dimethyl-butyl)-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-(3-Cyano-1-isopropyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-isobutyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 *N*-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 *N*-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-methyl-piperidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-propyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30

N-(3-Cyano-1-cyclopentyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 *N*-[3-Cyano-1-(1*H*-indol-2-ylmethyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-cycloheptyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-[3-Cyano-1-(1,2,3,4-tetrahydro-naphthalen-1-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(1-Bicyclo[2.2.1]hept-2-yl-3-cyano-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 *N*-[3-Cyano-1-(tetrahydro-pyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 *N*-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[3-Cyano-1-(tetrahydro-thiopyran-4-yl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 (4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

2-Bicyclo[4.1.0]hept-7-ylmethyl-*N*-(3-cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydro-naphthalen-2-ylmethyl)-butyramide;

5 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-indan-2-ylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclopentylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-{Cyano-[(cyclohexyl-ethyl-amino)-methyl]-methyl}-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(Cyano-piperidin-1-ylmethyl-methyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 *N*-[(3*S*)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 *N*-[(3*S*)-3-Cyano-1-cis-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3*S*)-3-Cyano-1-trans-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*-[(3*S*)-3-Cyano-1-cis-(4-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 *N*-[1-trans-(4-tert-Butyl-cyclohexyl)-(3*S*)-3--cyano-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-cyano-1-cis-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3S)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3S)-3-Cyano-1-cis-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3R)-3-Cyano-1-trans-(3-isopropyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 N-[(3S)-3-Cyano-1-cis-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

30 N-[(3S)-3-Cyano-1-trans-(2-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-cis-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 N-[(3S)-3-Cyano-1-trans-(2-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(2,2-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 N-[(3S)-3-Cyano-1-(4,4-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 N-[(3R)-3-Cyano-1-(2-methyl-2-phenyl-propyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

20 N-[(3S)-3-Cyano-1-indan-2-ylmethyl-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3S)-3-Cyano-1-(5-methyl-thiophen-2-ylmethyl)-pyrrolidin-3-yl]-(2S)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide and

25 the pharmaceutically acceptable salts, esters or tautomers thereof.

24. The compound according to claim 23 wherein the compound is selected from the group consisting of:

30

N-(cyano-1-benzyl-pyrrolidin-3-ylcarbamoyl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

5 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-(3-Cyano-1-cyclopropylmethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

10 *N*-(3-Cyano-1-phenethyl-pyrrolidin-3-yl)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[3-Cyano-1-(4-methyl-cyclohexyl)-pyrrolidin-3-yl]-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

15 (4*R*)-*N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-2-(4-methyl-cyclohexylmethyl)-4-morpholin-4-yl-4-oxo-butyramide;

20 *N*-(3-Cyano-1-cyclohexyl-pyrrolidin-3-yl)-4-morpholin-4-yl-4-oxo-2-(1,2,3,4-tetrahydro-naphthalen-2-ylmethyl)-butyramide;

N-[(3*S*)-3-Cyano-1-trans-(4-ethyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

25 *N*-[(3*S*)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide;

N-[(3*R*)-3-cyano-1-trans-(3-methyl-cyclohexyl)-pyrrolidin-3-yl]-(2*R*)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide and

30


N-[(3S)-3-Cyano-1-(3,3-dimethyl-cyclohexyl)-pyrrolidin-3-yl]-(2R)-2-cyclohexylmethyl-4-morpholin-4-yl-4-oxo-butyramide.

- 5 25. A pharmaceutical composition comprising a pharmaceutically effective amount of
a compound according to claim 1 or 11.
26. A method of modulating an autoimmune disease, said method comprising
administering to a patient in need of such treatment a pharmaceutically effect amount of a
10 compound according to claim 1 or 11.
27. The method according to claim 26 wherein the autoimmune disease is selected
from the group consisting of: rheumatoid arthritis, systemic lupus erythematosus, Crohn's
disease, ulcerative colitis, multiple sclerosis, Guillain-Barre syndrome, psoriasis, Grave's
15 disease, myasthenia gravis, scleroderma, glomerulonephritis, atopic dermatitis and
insulin-dependent diabetes mellitus.
28. A method of treating Alzheimer's disease comprising administering to a patient in
need of such treatment a pharmaceutically effective amount of a compound according to
20 claim 1 or 11.
29. A method of treating atherosclerosis comprising administering to a patient in
need of such treatment a pharmaceutically effective amount of a compound according to
claim 1 or 11.

30. A method of making a compound of the formula (Ia) according to claim 11


comprising:

- a) reacting a carboxylic acid compound shown below with an amine R_1H in the presence
 5 of a coupling reagent selected from the group consisting of EDC and HOBT, in a suitable solvent;



10


- b) hydrolyzing the product of step a) with aqueous base or acid to produce:



20

- c) reacting the product of step b) with an amino nitrile compound shown below in the presence of EDC as a coupling reagent, in a suitable solvent,

25




OR


30

d) reacting the product of step b) with an amino amide compound shown below in the presence of EDC as a coupling reagent, in a suitable solvent followed by dehydration of the primary amide to yield a nitrile group

5



10



15

wherein X is O, R₁ is an amino compound forming an amide bond with A, and R₂, R₃, R₄, R₅, R₆, R₇, and A are as defined in claim 11.


20

31. A method of making a compound of the formula(Ia) according to claim 11


comprising:

a) reacting an activated amide compound shown below with an aryl or heteroaryl organometallic reagent :


25




wherein R' is



b) hydrolyzing the product of step a) with aqueous base or acid to produce:




c) reacting the product of step b) with an amino nitrile compound shown below in the presence of EDC as a coupling reagent, in a suitable solvent,



or

d) reacting the product of step b) with an amino amide compound shown below in the presence of EDC as a coupling reagent, in a suitable solvent, followed by dehydration of the primary amide to yield a nitrile group,



- 15 and optionally reducing the carbonyl group attached to R1, to produce a compound of the formula (Ia) wherein X is O, R1 is alkyl, cycloalkyl, aryl or heteroaryl and R2, R3, R4, R5, R6, R7 and A are as defined in claim 11.

INTERNATIONAL SEARCH REPORT

Intern. Application No.

PCT/US 00/20453

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D295/18 C07D235/18 C07D211/66 C07D207/16 C07D403/06
 A61K31/5375 A61P37/00 C07C255/31

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, BEILSTEIN Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 24460 A (NOVARTIS ERFIND VERWALT GMBH ;ALTMANN EVA (CH); LATTMANN RENE (CH)) 20 May 1999 (1999-05-20) cited in the application claims	1-31
P,X	WO 99 56765 A (WARNER-LAMBERT COMPANY, USA; BASF AKTIENGESELLSCHAFT) 11 November 1999 (1999-11-11) examples 64,65	1-5, 11-13, 25,28



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

9 January 2001

Date of mailing of the international search report

17/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Pauwels, G

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/20453

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9924460 A	20-05-1999	AU 1487399 A	31-05-1999
		BR 9813197 A	29-08-2000
		EP 1028942 A	23-08-2000
		NO 20002320 A	04-07-2000
		ZA 9810073 A	05-05-1999
WO 9956765 A	11-11-1999	AU 3673099 A	23-11-1999

THIS PAGE BLANK (USPTO)